


Integrin cell surface interactions



European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://reactome.org/faq).

This is just an excerpt of a full-length report for this pathway. To access the complete report, please download it at the [Reactome Textbook](https://reactome.org/Textbook).

06/05/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)


Reactome database release: 88

This document contains 1 pathway and 37 reactions ([see Table of Contents](#))

Integrin cell surface interactions ↗

Stable identifier: R-BTA-216083

Inferred from: [Integrin cell surface interactions \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Collagen types III, IV, V, VI, VIII, IX, XVI bind integrins alpha1beta1 and alpha2beta1




Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-2327695

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [Collagen types III, IV, V, VI, VIII, IX, XVI bind integrins alpha1beta1 and alpha2beta1 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Collagen type I binds integrin alpha11beta1 ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-216045

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [Collagen type I binds integrin alpha11beta1 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Collagen type II binds integrin alpha10beta1 ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-216043

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [Collagen type II binds integrin alpha10beta1 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Collagen type XIII binds Integrin alpha1beta1 ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-2484965

Type: binding

Compartments: plasma membrane

Inferred from: [Collagen type XIII binds Integrin alpha1beta1 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Collagen types VI, IX bind integrin alpha10beta1 ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-4084903

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [Collagen types VI, IX bind integrin alpha10beta1 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Arresten binds integrin alpha1beta1 ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-4084912

Type: binding

Compartments: plasma membrane

Inferred from: [Arresten binds integrin alpha1beta1 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Endostatin binds integrin alphaVbeta3, alphaVbeta5, alpha3beta1, alpha5beta1 ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-4088264

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [Endostatin binds integrin alphaVbeta3, alphaVbeta5, alpha3beta1, alpha5beta1 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Tumstatin binds integrin alphaVbeta 3, alpha3beta1 ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-4085083

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [Tumstatin binds integrin alphaVbeta 3, alpha3beta1 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Canstatin binds integrins alphaVbeta3, alphaVbeta5 ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-4085087

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [Canstatin binds integrins alphaVbeta3, alphaVbeta5 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

THBS1 (Thrombospondin-1) binds Integrin alpha3beta1, alpha4beta1 ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-265429

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [THBS1 \(Thrombospondin-1\) binds Integrin alpha3beta1, alpha4beta1 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Integrins alpha4beta1, alpha8beta1, alphaVbeta1, alphaVbeta3, alphaVbeta6 bind Fibronectin matrix ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-216050

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [Integrins alpha4beta1, alpha8beta1, alphaVbeta1, alphaVbeta3, alphaVbeta6 bind Fibronectin matrix \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

SPP1 (osteopontin) binds integrin alpha5beta1, alpha9beta1 ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-265424

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [SPP1 \(osteopontin\) binds integrin alpha5beta1, alpha9beta1 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

SPP1 (osteopontin) binds CD44 ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-2752115

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [SPP1 \(osteopontin\) binds CD44 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

VCAM1 binds Integrin alpha4beta1 ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-198941

Type: binding

Compartments: plasma membrane

Inferred from: [VCAM1 binds Integrin alpha4beta1 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](http://www.pantherdb.org/about.jsp) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Integrin alpha4beta1 binds JAM2:JAM3 ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-202706

Type: binding

Compartments: plasma membrane

Inferred from: [Integrin alpha4beta1 binds JAM2:JAM3 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](http://www.pantherdb.org/about.jsp) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

ITGA4:ITGB1 binds natalizumab ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-9679740

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [ITGA4:ITGB1 binds natalizumab \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

MADCAM1-1 binds Integrin alpha4beta7 [↗](#)


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-199032

Type: binding

Compartments: plasma membrane

Inferred from: [MADCAM1-1 binds Integrin alpha4beta7 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Interaction of integrin alpha8beta1 with Tenascin-C ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-216064

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [Interaction of integrin alpha8beta1 with Tenascin-C \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Interaction of integrin alpha9beta1 with VCAM1 ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-265428

Type: binding

Compartments: plasma membrane

Inferred from: [Interaction of integrin alpha9beta1 with VCAM1 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Interaction of integrin alpha9beta1 with Tenascin-C ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-216068

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [Interaction of integrin alpha9beta1 with Tenascin-C \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

VTN (vitronectin) binds Integrin alphaVbeta3, alphaVbeta5, alphaVbeta8 ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-216076

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [VTN \(vitronectin\) binds Integrin alphaVbeta3, alphaVbeta5, alphaVbeta8 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Integrin alphaXbeta2 binds JAM3 ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-202704

Type: binding

Compartments: plasma membrane

Inferred from: [Integrin alphaXbeta2 binds JAM3 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

ICAM1-5 bind Integrin alphaLbeta2 (LFA-1) ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-199050

Type: binding

Compartments: plasma membrane

Inferred from: [ICAM1-5 bind Integrin alphaLbeta2 \(LFA-1\) \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Integrin alphaLbeta2 (LFA-1) binds F11R (JAM-A) ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-202718

Type: binding

Compartments: plasma membrane

Inferred from: [Integrin alphaLbeta2 \(LFA-1\) binds F11R \(JAM-A\) \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Integrin alphaMbeta2 (MAC1) binds JAM3 ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-202727

Type: binding

Compartments: plasma membrane

Inferred from: [Integrin alphaMbeta2 \(MAC1\) binds JAM3 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

BSG (basigin) binds Integrin alpha3beta1, alpha6beta1 ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-204434

Type: binding

Compartments: plasma membrane

Inferred from: [BSG \(basigin\) binds Integrin alpha3beta1, alpha6beta1 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Interaction of integrin alphaVbeta3 with IPSP (Bone sialoprotein 2) ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-265427

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [Interaction of integrin alphaVbeta3 with IPSP \(Bone sialoprotein 2\) \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Interaction of integrin alphaVbeta3 with Tenascin [↗](#)


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-265426

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [Interaction of integrin alphaVbeta3 with Tenascin \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Interaction of integrin alphaVbeta3 with Fibrillin [↗](#)


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-265423

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [Interaction of integrin alphaVbeta3 with Fibrillin \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Interaction of integrin alphaVbeta3 with von Willbrand Factor ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-265425

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [Interaction of integrin alphaVbeta3 with von Willbrand Factor \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Interaction of integrin alphaVbeta3 with PECAM1 ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-210304

Type: binding

Compartments: plasma membrane

Inferred from: [Interaction of integrin alphaVbeta3 with PECAM1 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Interaction of integrin alphaEbeta7 with Cadherin-1 ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-265422

Type: binding

Compartments: plasma membrane

Inferred from: [Interaction of integrin alphaEbeta7 with Cadherin-1 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Interaction of integrin alphaIIbeta3 with Fibronectin ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-349593

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [Interaction of integrin alphaIIbeta3 with Fibronectin \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Interaction of integrin alphaIIb beta3 with von Willebrand factor ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-216072

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [Interaction of integrin alphaIIb beta3 with von Willebrand factor \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Interaction of integrin alphaIIb beta 3 with THBS1 (Thrombospondin-1) ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-349603

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [Interaction of integrin alphaIIb beta 3 with THBS1 \(Thrombospondin-1\) \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

LUM (lumican) binds integrin alpha2beta1 ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-4085133

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [LUM \(lumican\) binds integrin alpha2beta1 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

COMP binds Integrin alpha5beta1, Integrin alphaVbeta3, CD47 ↗


Location: [Integrin cell surface interactions](#)

Stable identifier: R-BTA-2426259

Type: binding

Compartments: plasma membrane, extracellular region

Inferred from: [COMP binds Integrin alpha5beta1, Integrin alphaVbeta3, CD47 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Table of Contents

Introduction	1
🔧 Integrin cell surface interactions	2
🔗 Collagen types III, IV, V, VI, VIII, IX, XVI bind integrins alpha1beta1 and alpha2beta1	3
🔗 Collagen type I binds integrin alpha11beta1	4
🔗 Collagen type II binds integrin alpha10beta1	5
🔗 Collagen type XIII binds Integrin alpha1beta1	6
🔗 Collagen types VI, IX bind integrin alpha10beta1	7
🔗 Arresten binds integrin alpha1beta1	8
🔗 Endostatin binds integrin alphaVbeta3, alphaVbeta5, alpha3beta1, alpha5beta1	9
🔗 Tumstatin binds integrin alphaVbeta 3, alpha3beta1	10
🔗 Canstatin binds integrins alphaVbeta3, alphaVbeta5	11
🔗 THBS1 (Thrombospondin-1) binds Integrin alpha3beta1, alpha4beta1	12
🔗 Integrins alpha4beta1, alpha8beta1, alphaVbeta1, alphaVbeta3, alphaVbeta6 bind Fibronectin matrix	13
🔗 SPP1 (osteopontin) binds integrin alpha5beta1, alpha9beta1	14
🔗 SPP1 (osteopontin) binds CD44	15
🔗 VCAM1 binds Integrin alpha4beta1	16
🔗 Integrin alpha4beta1 binds JAM2:JAM3	17
🔗 ITGA4:ITGB1 binds natalizumab	18
🔗 MADCAM1-1 binds Integrin alpha4beta7	19
🔗 Interaction of integrin alpha8beta1 with Tenascin-C	20
🔗 Interaction of integrin alpha9beta1 with VCAM1	21
🔗 Interaction of integrin alpha9beta1 with Tenascin-C	22
🔗 VTN (vitronectin) binds Integrin alphaVbeta3, alphaVbeta5, alphaVbeta8	23
🔗 Integrin alphaXbeta2 binds JAM3	24
🔗 ICAM1-5 bind Integrin alphaLbeta2 (LFA-1)	25
🔗 Integrin alphaLbeta2 (LFA-1) binds F11R (JAM-A)	26
🔗 Integrin alphaMbeta2 (MAC1) binds JAM3	27
🔗 BSG (basigin) binds Integrin alpha3beta1, alpha6beta1	28
🔗 Interaction of integrin alphaVbeta3 with IPSP (Bone sialoprotein 2)	29
🔗 Interaction of integrin alphaVbeta3 with Tenascin	30
🔗 Interaction of integrin alphaVbeta3 with Fibrillin	31
🔗 Interaction of integrin alphaVbeta3 with von Willbrand Factor	32
🔗 Interaction of integrin alphaVbeta3 with PECAM1	33
🔗 Interaction of integrin alphaEbeta7 with Cadherin-1	34

↳ Interaction of integrin alphaIIb beta3 with Fibronectin	35
↳ Interaction of integrin alphaIIb beta3 with von Willebrand factor	36
↳ Interaction of integrin alphaIIb beta 3 with THBS1 (Thrombospondin-1)	37
↳ LUM (lumican) binds integrin alpha2beta1	38
↳ COMP binds Integrin alpha5beta1, Integrin alphaVbeta3, CD47	39
Table of Contents	40