


Factors involved in megakaryocyte development and platelet production



European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://reactome.org/licenses/).

This is just an excerpt of a full-length report for this pathway. To access the complete report, please download it at the [Reactome Textbook](https://reactome.org/textbook/).

06/05/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)


Reactome database release: 88

This document contains 2 pathways and 9 reactions ([see Table of Contents](#))

Factors involved in megakaryocyte development and platelet production ↗

Stable identifier: R-CEL-983231

Inferred from: Factors involved in megakaryocyte development and platelet production (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](http://www.pantherdb.org/about.jsp) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Dual-specific AKAPs bind type I and II PKA regulatory subunits ↗


Location: [Factors involved in megakaryocyte development and platelet production](#)

Stable identifier: R-CEL-992708

Type: binding

Compartments: cytosol

Inferred from: [Dual-specific AKAPs bind type I and II PKA regulatory subunits \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Mitofusins trans-interact linking mitochondria prior to fusion ↗


Location: Factors involved in megakaryocyte development and platelet production

Stable identifier: R-CEL-992703

Type: binding

Compartments: mitochondrial outer membrane

Inferred from: Mitofusins trans-interact linking mitochondria prior to fusion (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](http://www.pantherdb.org/about.jsp) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

HP1 alpha binds Histone H3K9(me)3 [↗](#)


Location: [Factors involved in megakaryocyte development and platelet production](#)

Stable identifier: R-CEL-994106

Type: binding

Compartments: nucleoplasm

Inferred from: [HP1 alpha binds Histone H3K9\(me\)3 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

LRRC16A binds F-actin capping protein ↗


Location: [Factors involved in megakaryocyte development and platelet production](#)

Stable identifier: R-CEL-994148

Type: binding

Compartments: cytosol

Inferred from: [LRRC16A binds F-actin capping protein \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: [F-actin capping protein is a heterodimer](#)

F-actin capping protein is a heterodimer ↗


Location: Factors involved in megakaryocyte development and platelet production

Stable identifier: R-CEL-879459

Type: binding

Compartments: cytosol

Inferred from: F-actin capping protein is a heterodimer (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](http://www.pantherdb.org/about.jsp) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Followed by: [LRR16A binds F-actin capping protein](#)

MICAL1 produces NADP+, H2O2 ↗


Location: [Factors involved in megakaryocyte development and platelet production](#)

Stable identifier: R-CEL-8865107

Type: omitted

Compartments: cytosol

Inferred from: [MICAL1 produces NADP+, H2O2 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

EHD proteins interact with Rabenosyn-5 ↗


Location: Factors involved in megakaryocyte development and platelet production

Stable identifier: R-CEL-1011576

Type: binding

Compartments: endosome membrane

Inferred from: EHD proteins interact with Rabenosyn-5 (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Followed by: Rabenosyn-5 connects Rab5 to VPS-45

Rabenosyn-5 connects Rab5 to VPS-45 ↗


Location: [Factors involved in megakaryocyte development and platelet production](#)

Stable identifier: R-CEL-1011600

Type: binding

Compartments: endosome membrane

Inferred from: [Rabenosyn-5 connects Rab5 to VPS-45 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: [EHD proteins interact with Rabenosyn-5](#)

DOCKs bind to RhoGEFs ↗


Location: Factors involved in megakaryocyte development and platelet production

Stable identifier: R-CEL-1011598

Type: binding

Compartments: cytosol

Inferred from: [DOCKs bind to RhoGEFs \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>


Kinesins ↗

Location: Factors involved in megakaryocyte development and platelet production

Stable identifier: R-CEL-983189

Compartments: cytosol

Inferred from: Kinesins (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](http://www.pantherdb.org/about.jsp) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Table of Contents

Introduction	1
☒ Factors involved in megakaryocyte development and platelet production	2
↳ Dual-specific AKAPs bind type I and II PKA regulatory subunits	3
↳ Mitofusins trans-interact linking mitochondria prior to fusion	4
↳ HP1 alpha binds Histone H3K9(me)3	5
↳ LRRRC16A binds F-actin capping protein	6
↳ F-actin capping protein is a heterodimer	7
☒ MICAL1 produces NADP+, H2O2	8
↳ EHD proteins interact with Rabenosyn-5	9
↳ Rabenosyn-5 connects Rab5 to VPS-45	10
↳ DOCKs bind to RhoGEFs	11
☒ Kinesins	12
Table of Contents	13