


MTOR signalling



European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://reactome.org/faq-fair-use/).

This is just an excerpt of a full-length report for this pathway. To access the complete report, please download it at the [Reactome Textbook](https://reactome.org/Textbook/).

29/04/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)


Reactome database release: 88

This document contains 4 pathways and 11 reactions ([see Table of Contents](#))

MTOR signalling ↗

Stable identifier: R-DME-165159

Inferred from: MTOR signalling (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.


[More details and caveats of the event inference in Reactome.](http://www.reactome.org) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Inhibition of TSC complex formation by PKB ↗

Location: MTOR signalling

Stable identifier: R-DME-165181

Inferred from: Inhibition of TSC complex formation by PKB (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](http://www.pantherdb.org/about.jsp) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

GTP loading by Rheb ↗


Location: [MTOR signalling](#)

Stable identifier: R-DME-165195

Type: transition

Compartments: cytosol, lysosomal membrane

Inferred from: [GTP loading by Rheb \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Followed by: [Formation of active mTORC1 complex](#)

Rag dimer formation ↗


Location: [MTOR signalling](#)

Stable identifier: R-DME-5653957

Type: binding

Compartments: cytosol

Inferred from: [Rag dimer formation \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Followed by: [Regulator binds Rag dimers](#)

Formation of Ragulator complex ↗


Location: [MTOR signalling](#)

Stable identifier: R-DME-5653936

Type: binding

Compartments: lysosomal membrane

Inferred from: [Formation of Ragulator complex \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Followed by: [Ragulator binds Rag dimers](#)

Ragulator binds Rag dimers ↗


Location: [MTOR signalling](#)

Stable identifier: R-DME-5653974

Type: binding

Compartments: cytosol, lysosomal membrane

Inferred from: [Ragulator binds Rag dimers \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: [Formation of Ragulator complex](#), [Rag dimer formation](#)

SLC38A9 transports L-Arg from lysosomal lumen to cytosol ↗


Location: [MTOR signalling](#)

Stable identifier: R-DME-8952726

Type: transition

Compartments: lysosomal lumen, cytosol, lysosomal membrane

Inferred from: [SLC38A9 transports L-Arg from lysosomal lumen to cytosol \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Ragulator:Rag dimers:SLC38A9 bind mTORC1 ↗


Location: [MTOR signalling](#)

Stable identifier: R-DME-5653968

Type: binding

Compartments: cytosol, lysosomal membrane

Inferred from: [Ragulator:Rag dimers:SLC38A9 bind mTORC1 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Formation of active mTORC1 complex ↗


Location: [MTOR signalling](#)

Stable identifier: R-DME-165680

Type: binding

Compartments: cytosol, lysosomal membrane

Inferred from: [Formation of active mTORC1 complex \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: [GTP loading by Rheb](#)

Followed by: [AKT1S1 \(PRAS40\) binds mTORC1](#)

AKT1S1 (PRAS40) binds mTORC1 ↗


Location: [MTOR signalling](#)

Stable identifier: R-DME-5672843

Type: binding

Compartments: cytosol, lysosomal membrane

Inferred from: [AKT1S1 \(PRAS40\) binds mTORC1 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: [Formation of active mTORC1 complex](#)

Followed by: [Activated Akt1 phosphorylates AKT1S1 \(PRAS40\)](#)

Activated Akt1 phosphorylates AKT1S1 (PRAS40) ↗


Location: [MTOR signalling](#)

Stable identifier: R-DME-377186

Type: transition

Compartments: cytosol, lysosomal membrane

Inferred from: [Activated Akt1 phosphorylates AKT1S1 \(PRAS40\) \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: [AKT1S1 \(PRAS40\) binds mTORC1](#)

Phosphorylated AKT1S1:mTORC1 binds YWHAB ↗


Location: [MTOR signalling](#)

Stable identifier: R-DME-5672824

Type: binding

Compartments: cytosol, lysosomal membrane

Inferred from: [Phosphorylated AKT1S1:mTORC1 binds YWHAB \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

mTORC1 phosphorylates AKT1S1 ↗


Location: [MTOR signalling](#)

Stable identifier: R-DME-5672828

Type: transition

Compartments: cytosol, lysosomal membrane

Inferred from: [mTORC1 phosphorylates AKT1S1 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.


[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

mTORC1-mediated signalling ↗

Location: MTOR signalling

Stable identifier: R-DME-166208

Inferred from: mTORC1-mediated signalling (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.


[More details and caveats of the event inference in Reactome.](http://www.pantherdb.org/about.jsp) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Energy dependent regulation of mTOR by LKB1-AMPK ↗

Location: [MTOR signalling](#)

Stable identifier: R-DME-380972

Inferred from: [Energy dependent regulation of mTOR by LKB1-AMPK \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Table of Contents

Introduction	1
❖ MTOR signalling	2
❖ Inhibition of TSC complex formation by PKB	3
➤ GTP loading by Rheb	4
➤ Rag dimer formation	5
➤ Formation of Ragulator complex	6
➤ Ragulator binds Rag dimers	7
➤ SLC38A9 transports L-Arg from lysosomal lumen to cytosol	8
➤ Ragulator:Rag dimers:SLC38A9 bind mTORC1	9
➤ Formation of active mTORC1 complex	10
➤ AKT1S1 (PRAS40) binds mTORC1	11
➤ Activated Akt1 phosphorylates AKT1S1 (PRAS40)	12
➤ Phosphorylated AKT1S1:mTORC1 binds YWHAB	13
➤ mTORC1 phosphorylates AKT1S1	14
❖ mTORC1-mediated signalling	15
❖ Energy dependent regulation of mTOR by LKB1-AMPK	16
Table of Contents	17