

# STAT92E is phosphorylated by phosphorylated HOP

Perrimon, N., Williams, MG.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://reactome.org/licenses/).

02/05/2024

## Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

## Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 88


This document contains 1 reaction ([see Table of Contents](#))

## STAT92E is phosphorylated by phosphorylated HOP [↗](#)

**Stable identifier:** R-DME-209210

**Type:** transition

**Compartments:** cytosol, plasma membrane



Tyr711 on STAT92E is phosphorylated by phosphorylated HOP tyrosine kinase.

### Literature references

- Small, S., Darnell JE, Jr., Desplan, C., Dearolf, CR., Yan, R. (1996). Identification of a Stat gene that functions in Drosophila development. *Cell*, 84, 421-30. [↗](#)
- Karsten, P., Perrimon, N., Zeidler, MP., Plischke, I. (2006). Mutational analysis reveals separable DNA binding and trans-activation of Drosophila STAT92E. *Cell Signal*, 18, 819-29. [↗](#)
- Nosaka, T., Ariyoshi, K., Miyajima, A., Onishi, M., Kitamura, T., Yamada, K. et al. (2000). Constitutive activation of STAT5 by a point mutation in the SH2 domain. *J Biol Chem*, 275, 24407-13. [↗](#)
- Chen, HW., Chen, X., Marinissen, MJ., Gutkind, JS., Oh, SW., Hou, SX. (2002). mom identifies a receptor for the Drosophila JAK/STAT signal transduction pathway and encodes a protein distantly related to the mammalian cytokine receptor family. *Genes Dev*, 16, 388-98. [↗](#)

### Editions

2006-11-02	Authored	Williams, MG.
2006-11-03	Edited	Williams, MG.
2008-01-16	Reviewed	Perrimon, N.