


Carnitine metabolism



European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://reactome.org/about/licenses).

This is just an excerpt of a full-length report for this pathway. To access the complete report, please download it at the [Reactome Textbook](https://reactome.org/textbook).

03/05/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 88


This document contains 1 pathway and 6 reactions ([see Table of Contents](#))

Carnitine metabolism ↗

Stable identifier: R-GGA-200425

Compartments: mitochondrion, cytosol

Inferred from: Carnitine metabolism (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](http://www.pantherdb.org/about.jsp) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

OCTN2 / SLC22A5 transports CAR from extracellular space to cytosol ↗


Location: [Carnitine metabolism](#)

Stable identifier: R-GGA-165026

Type: transition

Compartments: plasma membrane

Inferred from: [OCTN2 / SLC22A5 transports CAR from extracellular space to cytosol \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

CPT1A,B transfers PALM to CAR ↗


Location: Carnitine metabolism

Stable identifier: R-GGA-200406

Type: transition

Compartments: mitochondrial intermembrane space, cytosol, mitochondrial outer membrane

Inferred from: CPT1A,B transfers PALM to CAR (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](http://www.pantherdb.org/about.jsp) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Followed by: Exchange of palmitoylcarnitine and carnitine across the inner mitochondrial membrane

Exchange of palmitoylcarnitine and carnitine across the inner mitochondrial membrane ↗


Location: [Carnitine metabolism](#)

Stable identifier: R-GGA-200424

Type: transition

Compartments: mitochondrial inner membrane, mitochondrial intermembrane space, mitochondrial matrix

Inferred from: [Exchange of palmitoylcarnitine and carnitine across the inner mitochondrial membrane \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: [CPT1A,B transfers PALM to CAR](#)

Followed by: [palmitoylcarnitine + CoASH => palmitoyl-CoA + carnitine](#)

palmitoylcarnitine + CoASH => palmitoyl-CoA + carnitine ↗


Location: Carnitine metabolism

Stable identifier: R-GGA-200410

Type: transition

Compartments: mitochondrial inner membrane, mitochondrial matrix

Inferred from: palmitoylcarnitine + CoASH => palmitoyl-CoA + carnitine (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](http://www.pantherdb.org/about.jsp) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: Exchange of palmitoylcarnitine and carnitine across the inner mitochondrial membrane

MID1IP1 binds THRSP ↗


Location: [Carnitine metabolism](#)

Stable identifier: R-GGA-8866941

Type: binding

Compartments: cytosol

Inferred from: [MID1IP1 binds THRSP \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Btn-ACACA:2Mn2+ polymer carboxylates Ac-CoA to form Mal-CoA ↗


Location: Carnitine metabolism

Stable identifier: R-GGA-75851

Type: transition

Compartments: cytosol

Inferred from: Btn-ACACA:2Mn2+ polymer carboxylates Ac-CoA to form Mal-CoA (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](http://www.pantherdb.org/about.jsp) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Table of Contents

| | |
|--|---|
| Introduction | 1 |
| ☒ Carnitine metabolism | 2 |
| ↳ OCTN2 / SLC22A5 transports CAR from extracellular space to cytosol | 3 |
| ↳ CPT1A,B transfers PALM to CAR | 4 |
| ↳ Exchange of palmitoylcarnitine and carnitine across the inner mitochondrial membrane | 5 |
| ↳ palmitoylcarnitine + CoASH => palmitoyl-CoA + carnitine | 6 |
| ↳ MID1IP1 binds THRSP | 7 |
| ↳ Btn-ACACA:2Mn ²⁺ polymer carboxylates Ac-CoA to form Mal-CoA | 8 |
| Table of Contents | 9 |