

# Phosphorylation of ephexin1

Garapati, PV., Ip, NY.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of <u>Creative Commons Attribution 4.0 International (CC BY 4.0)</u> <u>License</u>. For more information see our <u>license</u>.

11/05/2024

# Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

### Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics, 18,* 142. 7
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. A
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res, 46*, D649-D655. ↗
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, *14*, e1005968. *オ*

This document contains 1 reaction (see Table of Contents)

## Phosphorylation of ephexin1 7

Stable identifier: R-GGA-3928596

#### Type: transition

#### Compartments: plasma membrane, cytosol



Activation of EPHAs leads to phosphorylation of ephexin1 (NGEF) on conserved tyrosine (Y) 179 (Y87 in isoform3) by Src family kinases (SFKs). This phosphorylation preferentially activates NGEF's GDP/GTP exchange activity specifically towards RHOA but not RAC1 and CDC42, thus switching the substrate preference of NGEF and leading to actin cytoskeletal changes that result in growth cone collapse (Sahin et al. 2005, Knoll and Drescher 2004).

#### Literature references

- Debant, A., Greenberg, ME., Corfas, G., Lin, MZ., Wright, TM., Schmidt, S. et al. (2005). Eph-dependent tyrosine phosphorylation of ephexin1 modulates growth cone collapse. *Neuron*, 46, 191-204.
- Knöll, B., Drescher, U. (2004). Src family kinases are involved in EphA receptor-mediated retinal axon guidance. J. Neurosci., 24, 6248-57. 🛪

#### **Editions**

2013-07-23	Authored, Edited	Garapati, P V.
2014-05-19	Reviewed	Ip, NY.