

# EHD proteins interact with Rabenosyn-5

Akkerman, JW., Jupe, S., Ouwehand, WH.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of <u>Creative Commons Attribution 4.0 International (CC BY 4.0)</u> <u>License</u>. For more information see our <u>license</u>.

25/09/2024

# Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

## Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics, 18,* 142. 7
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. A
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res, 46*, D649-D655.
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, *14*, e1005968. *对*

This document contains 1 reaction (see Table of Contents)

## EHD proteins interact with Rabenosyn-5 ↗

Stable identifier: R-HSA-1011576

Type: binding

#### Compartments: endosome membrane



The four human EH domain-containing proteins (EHD1-4) are a distinct highly-homologous subfamily of the Eps15homology (EH) domain family. They are distinct from most other EH family members in having the EH-domain at the C-terminus (Naslavsky & Caplan 2005). EH domains interact with other proteins; peptides containing Asp-Pro-Phe (NPF) motifs are major targets for EH-domain binding (Salcini et al. 1997). EH domain family proteins have regulatory roles in endocytic membrane transport events (Naslavsky & Caplan 2005); the EHD subfamily is believed to regulate endocytic recycling (George et al. 2007). All four human EHD proteins can rescue the vacuolated intestinal phenotype observed when the C. elegans orthologue rme-1 is mutated (George et al. 2007). Over 20 interaction partners have been reported for the C-terminal EHD proteins including clathirin, syndaptins and Arp2/3 (see Naslavsky & Caplan 2005). EHD1-3 all interact with Rabenosyn-5 (Rab5), a Rab5 effector (Naslavsky et al. 2004).

#### Literature references

Caplan, S., Naslavsky, N., Backlund PS, Jr., Boehm, M. (2004). Rabenosyn-5 and EHD1 interact and sequentially regulate protein recycling to the plasma membrane. *Mol Biol Cell, 15*, 2410-22.

#### Editions

2010-10-29	Reviewed	Akkerman, JW.
2010-11-12	Edited	Jupe, S.
2010-11-12	Authored	Ouwehand, WH.