

HNP1-3 bind gp120

Jupe, S., McDermott, AM.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://reactome.org/licenses/).

08/05/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 88


This document contains 1 reaction ([see Table of Contents](#))

HNP1-3 bind gp120 [↗](#)

Stable identifier: R-HSA-1471354

Type: binding

Compartments: extracellular region, viral envelope



Alpha-defensins, theta-defensins and their synthetic analogues the retrocyclins have been shown in numerous studies to have anti-HIV-1 activity (Chang & Klotman 2004). This appears to be mediated via multiple mechanisms including direct viral inactivation and down regulation of host-cell target co-receptors important for viral entry (Furci et al. 2007, Seidel et al. 2010). HNP1-3 act as lectins, binding with relatively high affinity to gp120 (KD range, 15.8-52.8 nM) on the HIV-1 envelope and CD4 (KD range, 8.0-34.9 nM) on host target cells, both important molecules for viral entry (Wang et al. 2004). Retrocyclins, artificial theta defensins predicted from human defensin pseudogenes, bind with even higher affinity whereas HNP-4 binding is much weaker (Wu et al. 2005). Alpha defensins have been demonstrated to inhibit the binding of gp120 to CD4 thus blocking HIV-1 fusion with its target cells (Furci et al. 2007).

Literature references

Waring, AJ., Lal, RB., Hong, T., Cole, AM., Owen, SM., Wang, W. et al. (2004). Activity of alpha- and theta-defensins against primary isolates of HIV-1. *J Immunol*, 173, 515-20. [↗](#)

Editions

2011-04-28	Authored	Jupe, S.
2011-07-27	Edited	Jupe, S.
2011-11-03	Reviewed	McDermott, AM.