

IRAK1 or IRAK2 binds to the activated IRAK4 :activated TLR:MyD88:TIRAP com- plex

Gay, NJ., Gillespie, ME., Napetschnig, J., Shamovsky, V., de Bono, B.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://reactome.org/licenses/).

29/04/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 88

This document contains 1 reaction ([see Table of Contents](#))


IRAK1 or IRAK2 binds to the activated IRAK4 :activated TLR:MyD88:TIRAP complex



Stable identifier: R-HSA-166091

Type: binding

Compartments: plasma membrane, cytosol



IRAK2 has been implicated in IL1R and TLR signaling by the observation that IRAK2 can associate with MyD88 and Mal (Muzio et al. 1997). Like IRAK1, IRAK2 is activated downstream of IRAK4 (Kawagoe et al. 2008). It has been suggested that IRAK1 activates IRAK2 (Wesche et al. 1999) but IRAK2 phosphorylation is observed in IRAK1^{-/-} mouse macrophages while IRAK4 deficiency abrogates IRAK2 phosphorylation (Kawagoe et al. 2008), suggesting that activated IRAK4 phosphorylates IRAK2 as it does IRAK1. IL6 production in response to IL1beta is impaired in embryonic fibroblasts from IRAK1 or IRAK2 knockout mice and abrogated in IRAK1/2 dual knockouts (Kawagoe et al. 2007) suggesting that IRAK1 and IRAK2 are both involved in IL1R signaling downstream of IRAK4.

IRAK1 is recruited to the TLR complex through binding with IRAK4[Lin SC et al 2010].

Literature references

- Szymak, J., Keating, SE., Bowie, AG., Flannery, SM. (2011). Human interleukin-1 receptor-associated kinase-2 is essential for Toll-like receptor-mediated transcriptional and post-transcriptional regulation of tumor necrosis factor alpha. *J. Biol. Chem.*, 286, 23688-97. [↗](#)
- Maloney, GM., Bowie, AG., Keating, SE., Moran, EM. (2007). IRAK-2 participates in multiple toll-like receptor signaling pathways to NFkappaB via activation of TRAF6 ubiquitination. *J Biol Chem*, 282, 33435-43. [↗](#)
- Rao, NL., Gottipati, S., Fung-Leung, WP. (2008). IRAK1: a critical signaling mediator of innate immunity. *Cell Signal*, 20, 269-76. [↗](#)
- Lo, YC., Lin, SC., Wu, H. (2010). Helical assembly in the MyD88-IRAK4-IRAK2 complex in TLR/IL-1R signalling. *Nature*, 465, 885-90. [↗](#)
- Li, X., Wan, Y., Stark, GR., Thomas, J., Hamilton, T., Kim, TW. et al. (2009). Interleukin-1 receptor-associated kinase 2 is critical for lipopolysaccharide-mediated post-transcriptional control. *J Biol Chem*, 284, 10367-75. [↗](#)

Editions

2005-08-16	Authored	de Bono, B.
2006-04-24	Reviewed	Gay, NJ.
2010-08-25	Revised	Shamovsky, V.
2010-11-30	Reviewed	Gillespie, ME.
2012-11-06	Edited	Shamovsky, V.
2012-11-16	Reviewed	Napetschnig, J.