

IKKbeta is activated

Annibali, D., Chao, MV., Friedman, WJ., Jassal, B., Nasi, S.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](#). For more information see our [license](#).

08/05/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 88


This document contains 1 reaction ([see Table of Contents](#))

IKKbeta is activated ↗

Stable identifier: R-HSA-193703

Type: transition

Compartments: cytosol, plasma membrane



Atypical PKC isoforms phosphorylate the beta subunit of the IKK complex (on Serines 177 and 181) thereby serving as an IKK kinase. TRAF6 and p62 as well appear to have a role in IKK activation. TRAF6 mediates the assembly of K63-linked poly-Ub chains required for IKK activation. The ubiquitin binding property of p62 may also be relevant in regulating IKK activation.

Literature references

Bren, G., Moscat, J., Diaz-Meco, MT., Paya, CV., Lallena, MJ. (1999). Activation of IkappaB kinase beta by protein kinase C isoforms. *Mol Cell Biol*, 19, 2180-8. ↗

Editions

2006-10-10	Authored	Annibali, D., Nasi, S.
2008-05-20	Reviewed	Friedman, WJ.
2008-05-20	Edited	Jassal, B.
2008-05-28	Reviewed	Chao, MV.