

MHC class II antigen internalization

Garapati, PV., Neefjes, J.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of <u>Creative Commons Attribution 4.0 International (CC BY 4.0)</u> <u>License</u>. For more information see our <u>license</u>.

20/03/2025

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Fabregat, A., Fabregat, A., Fabregat, A., Fabregat, A., Fabregat, A. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. 7
- Sidiropoulos, K., Sidiropoulos, K., Sidiropoulos, K., Sidiropoulos, K., Sidiropoulos, K., Sidiropoulos, K. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467.
- Fabregat, A., Fabregat, A., Fabregat, A., Fabregat, A., Fabregat, A., Fabregat, A. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res, 46*, D649-D655.
- Fabregat, A., Fabregat, A., Fabregat, A., Fabregat, A., Fabregat, A., Fabregat, A. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, *14*, e1005968. *¬*

This document contains 1 reaction (see Table of Contents)

MHC class II antigen internalization 7

Stable identifier: R-HSA-2130627

Type: omitted

Compartments: extracellular region, lysosomal lumen



MHC II typically presents antigens derived from exogenous proteins internalized by APCs such as macrophages, B cells or dendritic cells. Different types of antigen use different routes of internalization. Endocytosis may be specific, mediated by a range of receptors expressed on APC, or may occur by nonspecific mechanisms such as phagocytosis, macropinocytosis or autophagy. Antigens are first loaded into endocytic vesicles and progress along the early endosomes (EE)-late endosomes (LE) lysosomal axis. Antigens are exposed to increasingly acidic, more denaturing and proteolytic conditions (Doherty & McMahon 2009, Underhill & Ozinsky 2002).

Literature references

Mellman, I. (1996). Endocytosis and molecular sorting. Annu Rev Cell Dev Biol, 12, 575-625. 🛪

- Harding, CV., Noss, E., Ramachandra, L., Boom, WH. (2001). Processing of Mycobacterium tuberculosis antigen 85B involves intraphagosomal formation of peptide-major histocompatibility complex II complexes and is inhibited by live bacilli that decrease phagosome maturation. *J Exp Med*, *194*, 1421-32. *¬*
- Harding, CV., Ramachandra, L., Song, R. (1999). Phagosomes are fully competent antigen-processing organelles that mediate the formation of peptide:class II MHC complexes. *J Immunol, 162,* 3263-72.

McMahon, HT., Doherty, GJ. (2009). Mechanisms of endocytosis. Annu Rev Biochem, 78, 857-902. 🛪

Editions

2012-02-21	Authored, Edited	Garapati, P V.
2012-05-14	Reviewed	Neefjes, J.