

# Translocation of TGN-lysosome vesicle to

## lysosome

Garapati, P V., Neefjes, J.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of <u>Creative Commons Attribution 4.0 International (CC BY 4.0)</u> <u>License</u>. For more information see our <u>license</u>.

03/05/2024

## Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

### Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics, 18,* 142. 7
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. A
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res, 46*, D649-D655. ↗
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, *14*, e1005968. *オ*


This document contains 1 reaction (see Table of Contents)

## Translocation of TGN-lysosome vesicle to lysosome

Stable identifier: R-HSA-2130641

#### Type: transition

#### Compartments: trans-Golgi network membrane, cytosol



Budding vesicles loaded with the nonameric complex bud off from the TGN to reach the late-endosome/lysosome membrane.

#### Literature references

- Peterson, PA., Tainer, JA., Pond, L., Schutze, MP., Jackson, MR., Teyton, L. et al. (1995). A role for acidic residues in di-leucine motif-based targeting to the endocytic pathway. J Biol Chem, 270, 19989-97. *¬*
- Davidson, HW. (1999). Direct transport of newly synthesized HLA-DR from the trans-Golgi network to major histocompatibility complex class II containing compartments (MIICS) demonstrated using a novel tyrosine-sulfated chimera. J Biol Chem, 274, 27315-22. ↗
- Goud, B., Hoflack, B., Salamero, J., Le Borgne, R., Saudrais, C. (1996). Expression of major histocompatibility complex class II molecules in HeLa cells promotes the recruitment of AP-1 Golgi-specific assembly proteins on Golgi membranes. J Biol Chem, 271, 30318-21. ↗
- Morelli, MA., Frank, RW., Saviano, G., Bremnes, B., Motta, A., Bakke, O. (1995). Structure-activity relationship of the leucine-based sorting motifs in the cytosolic tail of the major histocompatibility complex-associated invariant chain. J Biol Chem, 270, 27165-71.
- Rodionov, DG., Bakke, O. (1998). Medium chains of adaptor complexes AP-1 and AP-2 recognize leucine-based sorting signals from the invariant chain. *J Biol Chem*, 273, 6005-8.

### **Editions**

2012-02-21	Authored, Edited	Garapati, P V.
2012-05-14	Reviewed	Neefjes, J.