

PTK2 autophosphorylates at Y397

Garapati, P V., Ip, NY.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://reactome.org/licenses/).

13/05/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)

Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)

Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)

Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 88

This document contains 1 reaction ([see Table of Contents](#))


PTK2 autophosphorylates at Y397 ↗

Stable identifier: R-HSA-3928610

Type: transition

Compartments: plasma membrane, cytosol

Inferred from: [Ptk2 autophosphorylates at Y397 \(Mus musculus\)](#)



Focal adhesion kinase 1 (PTK2, FAK, FAK1) activation plays a critical role in EPHB receptor signaling in dendritic spines. PTK2 has six tyrosine phosphorylation sites, with tyrosine 397 being the main auto-phosphorylation site present upstream of the kinase domain (Schaller et al. 1994). Activation of EPHB receptors induces long-lasting phosphorylation of PTK2 on tyrosine 397 (Shi et al. 2009). This phosphorylated tyrosine then creates a binding site for other signaling proteins that link PTK2 to downstream signaling pathways and actin cytoskeleton.

Editions

2013-07-23	Authored, Edited	Garapati, P V.
2014-05-19	Reviewed	Ip, NY.