

# trans-Golgi Network Coat Assembly

Gillespie, ME., Simpson, JC.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of <u>Creative Commons Attribution 4.0 International (CC BY 4.0)</u> <u>License</u>. For more information see our <u>license</u>.

20/05/2024

# Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

## Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics, 18,* 142. 7
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. A
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res, 46*, D649-D655. ↗
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, *14*, e1005968. *オ*

This document contains 1 reaction (see Table of Contents)

## trans-Golgi Network Coat Assembly 7

#### Stable identifier: R-HSA-421831

Type: dissociation

#### Compartments: Golgi membrane, cytosol



Once the basic components of the docking complex are assembled with one end of AP-1 bound to cargo molecules, the other end binds to clathrin. Clathrin triskelions polymerize into hexagons and pentagons, forming a cage, which leads to membrane deformation. This polymerization step drives the sculpting of the vesicle. The number of clathrin triskelions required to sculpt a vesicle appears to be variable, but has been estimated to require 36 - 60 triskelions assocaited with 30 - 66 AP-1 complexes. Here a ~380 angstroms vesicle is represented with 48 clathrin triskelions and 52 AP-1 complexes.

#### Literature references

- James, DE., Bryant, NJ., Govers, R. (2002). Regulated transport of the glucose transporter GLUT4. *Nat Rev Mol Cell Biol*, *3*, 267-77. *¬*
- Roseman, AM., Pearse, BM., Smith, CJ., Musacchio, A., Kirchhausen, Tomas., Harrison, SC. (1999). Functional organization of clathrin in coats: combining electron cryomicroscopy and X-ray crystallography. *Mol Cell, 3*, 761-70.

#### **Editions**

2008-05-22	Edited	Gillespie, ME.
2009-08-27	Authored	Gillespie, ME.
2009-08-28	Reviewed	Simpson, JC.