


ERCC6 (CSB) and EHMT2 (G9a) positively regulate rRNA expression



Comai, L., Gillespie, ME., Iben, S., May, B., Shiao, YH.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://reactome.org/licenses/).

This is just an excerpt of a full-length report for this pathway. To access the complete report, please download it at the [Reactome Textbook](https://reactome.org/textbook/).

17/04/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)


Reactome database release: 88

This document contains 1 pathway and 4 reactions ([see Table of Contents](#))

ERCC6 (CSB) and EHMT2 (G9a) positively regulate rRNA expression ↗

Stable identifier: R-HSA-427389

Compartments: nucleoplasm



About half of the rRNA genes in the genome are actively expressed, being transcribed by RNA polymerase I (reviewed in Nemeth and Langst 2008, Bartova et al. 2010, Goodfellow and Zomerdijk 2012, Grummt and Langst 2013). As inferred from mouse, those genes that are expressed are activated by ERCC6 (also known as Cockayne Syndrome protein, CSB) which interacts with TTF-I bound to the T0 terminator region (also known as the Sal Box) of rRNA genes (Yuan et al. 2007, reviewed in Birch and Zomerdijk 2008, Grummt and Langst 2013). ERCC6 recruits the histone methyltransferase EHMT2 (also known as G9a) which dimethylates histone H3 at lysine-9 in the coding region of rRNA genes. The dimethylated lysine is bound by CBX3 (also known as Heterochromatic Protein-1gamma, HP1gamma) and increases expression of the rRNA gene. Continuing dimethylation depends on continuing transcription. Mutations in CSB result in dysregulation of RNA polymerase I transcription, which plays a role in the symptoms of Cockayne Syndrome (reviewed in Hannan et al. 2013).

Literature references

- Uhlířová, R., Raska, I., Kozubek, S., Galiová, G., Orlova, D., Bártová, E. et al. (2010). Structure and epigenetics of nucleoli in comparison with non-nucleolar compartments. *J. Histochem. Cytochem.*, 58, 391-403. ↗
- Zhou, Y., Grummt, I., Feng, W., Yuan, X., Imhof, A. (2007). Activation of RNA polymerase I transcription by cockayne syndrome group B protein and histone methyltransferase G9a. *Mol Cell*, 27, 585-95. ↗
- Zomerdijk, JC., Goodfellow, SJ. (2012). Basic mechanisms in RNA polymerase I transcription of the ribosomal RNA genes. *Subcell. Biochem.*, 61, 211-36. ↗
- Rothblum, LI., Pearson, RB., Sanij, E., Hannan, RD., Hannan, KM. (2013). Dysregulation of RNA polymerase I transcription during disease. *Biochim. Biophys. Acta*, 1829, 342-60. ↗
- Längst, G., Grummt, I. (2013). Epigenetic control of RNA polymerase I transcription in mammalian cells. *Biochim. Biophys. Acta*, 1829, 393-404. ↗

Editions

2009-06-21	Authored, Edited	May, B.
2016-02-12	Reviewed	Iben, S.

TTF-I binds to the Sal Box ↗


Location: ERCC6 (CSB) and EHMT2 (G9a) positively regulate rRNA expression

Stable identifier: R-HSA-74987

Type: binding

Compartments: nucleoplasm

Inferred from: Ttf-I binds the T0 region (Sal Box) of the rDNA (Mus musculus)



As inferred from mouse cell models, the Transcription termination factor (TTF1, also known as TTF-1 and TTF-I) binds an 18 base pair sequence element known as the Sal Box found in multiple copies in the nontranscribed spacer downstream of the 28S rRNA coding region. This element is the termination signal for ribosomal gene transcription. Binding of TTF1 mediates the pausing of the elongating transcription complex. TTF1 has a relatively low affinity for purified DNA but binds cooperatively to chromatin. Oligomers of TTF1 interact in trans to bind adjacent intergenic regions and form loops of the rDNA. Binding of TTF1 to the Sal Box is also influenced by interaction of TTF1 with TIP5 and possibly other proteins.

Followed by: Recruitment of ERCC6 (CSB), EHMT2 (G9a), and NuRD to the promoter of rRNA gene

Literature references

Evers, R., Grummt, I. (1995). Molecular coevolution of mammalian ribosomal gene terminator sequences and the transcription termination factor TTF-I. *Proc Natl Acad Sci U S A*, 92, 5827-31. ↗

Editions

2003-07-03	Authored	Comai, L.
2014-02-18	Reviewed	Shiao, YH.
2016-02-12	Reviewed	Iben, S.
2024-03-06	Edited	Gillespie, ME.

Recruitment of ERCC6 (CSB), EHMT2 (G9a), and NuRD to the promoter of rRNA gene

↗


Location: ERCC6 (CSB) and EHMT2 (G9a) positively regulate rRNA expression

Stable identifier: R-HSA-427404

Type: binding

Compartments: nucleoplasm

Inferred from: Recruitment of CSB, G9a, and NuRD to the rRNA promoter (Homo sapiens)



Transcription Termination Factor-I (TTF-I) is a sequence-specific binding protein that binds sites 5' (Tsp and T0 sites) and 3' (T1-10 site) of rRNA genes. As inferred from mouse, when TTF-I is bound to the promoter-proximal T0 site TTF-I either recruits ERCC6 (also known as Cockayne Syndrome Protein, CSB), EHMT2 (also known as histone methyltransferase G9a), and NuRD to activate expression (Shimono et al. 2005, Lebedev et al. 2008) or recruits the Nucleolar Remodeling Complex (NoRC) to repress expression. How one is selected over the other is unknown.

CHD4 and presumably the rest of the NuRD complex is associated with bivalent domains containing H3K4me3 (active chromatin mark) and H3K27me3 (inactive chromatin mark). ERCC6 and EHMT2 appear to cooperate to regulate activation of rRNA expression with ERCC6 mediating the transition to permissive chromatin (Lebedev et al. 2008) and EHMT2 mediating the transition to active chromatin, which involves the positional shift of one nucleosome at the promoter.

Preceded by: TTF-I binds to the Sal Box

Followed by: TTF1:rRNA promoter:ERCC6:EHMT2 complex dimethylates histone H3 at lysine-9.

Literature references

- Lebedev, A., Scharffetter-Kochanek, K., Iben, S. (2008). Truncated Cockayne syndrome B protein represses elongation by RNA polymerase I. *J Mol Biol*, 382, 266-74. ↗
- Shimono, Y., Shimono, K., Ishiguro, N., Takahashi, M., Shimokata, K. (2005). Microspherule protein 1, Mi-2beta, and RET finger protein associate in the nucleolus and up-regulate ribosomal gene transcription. *J. Biol. Chem.*, 280, 39436-47. ↗
- Vonesch, JL., Auriol, J., Egly, JM., Iben, S., Proietti de Santis, L., Bradsher, J. et al. (2002). CSB is a component of RNA pol I transcription. *Mol. Cell*, 10, 819-29. ↗

Editions

2009-06-19	Edited	May, B.
2009-06-21	Authored	May, B.
2016-02-12	Reviewed	Iben, S.

TTF1:rRNA promoter:ERCC6:EHMT2 complex dimethylates histone H3 at lysine-9. ↗


Location: ERCC6 (CSB) and EHMT2 (G9a) positively regulate rRNA expression

Stable identifier: R-HSA-427336

Type: transition

Compartments: nucleoplasm

Inferred from: Ttf-I:rRNA Promoter:Ercc6:Ehmt2 complex dimethylates histone H3 at lysine-9 (Homo sapiens)



As inferred from mouse, EHMT2 (histone methyltransferase G9a) dimethylates histone H3 at lysine-9 (H3K9me2) in the transcribed region of the rRNA gene. Dimethylation of histone H3 in the transcribed region causes increased rRNA expression, which contrasts with the repressive effect of H3K9me2 in other regions of the genome. The histone binding activity and ATPase activity of CHD4 in the NuRD complex are also needed for activation.

Preceded by: Recruitment of ERCC6 (CSB), EHMT2 (G9a), and NuRD to the promoter of rRNA gene

Followed by: CBX3 (HP1gamma) binds histone H3 dimethylated at lysine-9

Editions

2009-06-19	Edited	May, B.
2009-06-21	Authored	May, B.
2016-02-12	Reviewed	Iben, S.

CBX3 (HP1gamma) binds histone H3 dimethylated at lysine-9 ↗


Location: [ERCC6 \(CSB\) and EHMT2 \(G9a\) positively regulate rRNA expression](#)

Stable identifier: R-HSA-427383

Type: binding

Compartments: nucleoplasm

Inferred from: [Cbx3 \(HP1gamma\) Binds Histone H3 dimethylated at lysine-9 \(Mus musculus\)](#)



As inferred from mouse, CBX3 (Heterochromatic Protein 1gamma, HP1gamma) binds histone H3 dimethylated at lysine-9 (H3K9me2). In other regions of the genome, CBX3 can be associated with repression of transcription, however dimethylated H3 lysine-9 and CBX3 in the transcribed region of the rRNA gene are associated with enhanced expression. CBX3 bound to gene bodies can facilitate cotranscriptional processing of RNA (Smallwood et al. 2012).

Preceded by: [TTF1:rRNA promoter:ERCC6:EHMT2 complex dimethylates histone H3 at lysine-9.](#)

Literature references

Smallwood, A., Ren, B., Hon, GC., Jin, F., Henry, RE., Espinosa, JM. (2012). CBX3 regulates efficient RNA processing genome-wide. *Genome Res.*, 22, 1426-36. ↗

Editions

2009-06-19	Edited	May, B.
2009-06-21	Authored	May, B.
2016-02-12	Reviewed	Iben, S.

Table of Contents

Introduction	1
☒ ERCC6 (CSB) and EHMT2 (G9a) positively regulate rRNA expression	2
↳ TTF-I binds to the Sal Box	4
↳ Recruitment of ERCC6 (CSB), EHMT2 (G9a), and NuRD to the promoter of rRNA gene	5
↳ TTF1:rRNA promoter:ERCC6:EHMT2 complex dimethylates histone H3 at lysine-9.	6
↳ CBX3 (HP1gamma) binds histone H3 dimethylated at lysine-9	7
Table of Contents	8