

IGF2BP3 binds specific RNAs

Chao, JA., May, B., Singer, RH.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](#). For more information see our [license](#).

06/05/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 88


This document contains 1 reaction ([see Table of Contents](#))

IGF2BP3 binds specific RNAs [↗](#)

Stable identifier: R-HSA-428287

Type: binding

Compartments: cytosol



Insulin-like Growth Factor mRNA Binding Factor-3 (IGF2BP3, also known as IMP3 and VICKZ3) binds several specific RNAs containing the sequence motif CAUH (where H is A, C, or U). Binding causes stabilization and subcellular localization of the RNA.

Isoforms of IGF2 mRNA containing leader-3 are bound by IGF2BP3 at the 5' UTR, repressing translation (other isoforms of IGF2 are constitutive).

Literature references

Nielsen, J., Wewer, UM., Christiansen, J., Johnsen, AH., Lykke-Andersen, J., Nielsen, FC. (1999). A family of insulin-like growth factor II mRNA-binding proteins represses translation in late development. *Mol Cell Biol*, 19, 1262-70. [↗](#)

Nielsen, J., Christiansen, J., Nielsen, FC. (2001). A family of IGF-II mRNA binding proteins (IMP) involved in RNA trafficking. *Scand J Clin Lab Invest Suppl*, 234, 93-9. [↗](#)

Yisraeli, JK. (2005). VICKZ proteins: a multi-talented family of regulatory RNA-binding proteins. *Biol Cell*, 97, 87-96. [↗](#)

Wewer, UM., Hansen, TV., Christiansen, J., Borup, R., Jønson, L., Vikesaa, J. et al. (2006). RNA-binding IMPs promote cell adhesion and invadopodia formation. *EMBO J*, 25, 1456-68. [↗](#)

Landthaler, M., Tuschl, T., Ulrich, A., Ascano M, Jr., Hausser, J., Khorshid, M. et al. (2010). Transcriptome-wide identification of RNA-binding protein and microRNA target sites by PAR-CLIP. *Cell*, 141, 129-41. [↗](#)

Editions

2009-07-05	Authored, Edited	May, B.
2010-05-30	Reviewed	Singer, RH., Chao, JA.