

BH3-only proteins associate with and inac-

tivate anti-apoptotic BCL-2

Matthews, L.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of <u>Creative Commons Attribution 4.0 International (CC BY 4.0)</u> <u>License</u>. For more information see our <u>license</u>.

14/05/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics, 18,* 142. 7
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. A
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res, 46*, D649-D655. ↗
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, *14*, e1005968. *オ*


This document contains 1 reaction (see Table of Contents)

BH3-only proteins associate with and inactivate anti-apoptotic BCL-2 7

Stable identifier: R-HSA-508163

Type: binding

Compartments: mitochondrial outer membrane



Bcl-2 interacts with tBid (Yi et al. 2003), BIM (Puthalakath et al. 1999), PUMA (Nakano and Vousden 2001), NOXA (Oda et al. 2000), BAD (Yang et al. 2005), BMF (Puthalakath et al. 2001), resulting in inactivation of BCL2.

Literature references

- Beaumont, JG., O'Reilly, LA., Cheney, RE., Coultas, L., Villunger, A., Puthalakath, H. et al. (2001). Bmf: a proapoptotic BH3-only protein regulated by interaction with the myosin V actin motor complex, activated by anoikis. *Science* , 293, 1829-32. ↗
- Boise, LH., Jockel, J., Zha, J., Yang, E., Korsmeyer, SJ., Thompson, CB. (1995). Bad, a heterodimeric partner for Bcl-XL and Bcl-2, displaces Bax and promotes cell death. *Cell*, 80, 285-91. *¬*
- O'Reilly, LA., Puthalakath, H., Strasser, A., King, SM., Huang, DC. (1999). The proapoptotic activity of the Bcl-2 family member Bim is regulated by interaction with the dynein motor complex. *Mol Cell*, *3*, 287-96.
- Dong, Z., Yin, XM., Yi, X. (2003). Inhibition of Bid-induced apoptosis by Bcl-2. tBid insertion, Bax translocation, and Bax/Bak oligomerization suppressed. *J Biol Chem, 278*, 16992-9. 🛪
- Taniguchi, T., Tokino, T., Yamashita, T., Shibue, T., Ohki, R., Oda, E. et al. (2000). Noxa, a BH3-only member of the Bcl-2 family and candidate mediator of p53-induced apoptosis. *Science*, *288*, 1053-8. *¬*

Editions

2010-02-05

Authored

Matthews, L.