

# Defective POMT1 does not transfer Man from Dol-P-Man to DAG1

Hansen, L., Jassal, B., Joshi, HJ.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of <u>Creative Commons Attribution 4.0 International (CC BY 4.0)</u> <u>License</u>. For more information see our <u>license</u>.

19/05/2024

## Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

## Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics, 18*, 142. 7
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. A
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res, 46*, D649-D655. ↗
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, *14*, e1005968. *オ*

This document contains 1 reaction (see Table of Contents)

## Defective POMT1 does not transfer Man from Dol-P-Man to DAG1 7

Stable identifier: R-HSA-5615604

#### Type: transition

**Compartments:** endoplasmic reticulum membrane, cytosol, integral component of cytoplasmic side of endoplasmic reticulum membrane

#### Diseases: muscular dystrophy-dystroglycanopathy



Co-expression of both protein O-mannosyl-transferases 1 and 2 (POMT1 and POMT2; CAZy family GT39) is necessary for enzyme activity, that is mediating the transfer of mannosyl residues to the hydroxyl group of serine or threonine residues of proteins such as alpha-dystroglycan (DAG1; MIM:128239). DAG1 is a cell surface protein that plays an important role in the assembly of the extracellular matrix in muscle, brain, and peripheral nerves by linking the basal lamina to cytoskeletal proteins. Defects in POMT1 (MIM:607423) results in defective glycosylation of DAG1 and can cause severe congenital muscular dystrophy-dystroglycanopathies ranging from a severe type A, MDDGA1 (brain and eye abnormalities; MIM:236670), through a less severe type B, MDDGB1 (congenital form with mental retardation; MIM:613155) to a milder type C, MDDGC1 (limb girdle form; MIM:609308) (Bertini et al. 2011, Wells 2013).

Several mutations are known and mutations causing the severest type A1 form include G76R, Q303\*, S727Afs\*3, W705Lfs\*26 and D723Efs\*8 (Beltran-Valero de Bernabe et al. 2002, Godfrey et al. 2007, Mercuri et al. 2009).

#### Literature references

- Talim, B., Smith, J., Abbs, S., Brown, SC., Clement, E., Godfrey, C. et al. (2007). Refining genotype phenotype correlations in muscular dystrophies with defective glycosylation of dystroglycan. *Brain, 130*, 2725-35.
- Scuderi, C., Mottarelli, E., Morandi, L., Laverda, A., Saredi, S., Moggio, M. et al. (2009). Congenital muscular dystrophies with defective glycosylation of dystroglycan: a population study. *Neurology*, *72*, 1802-9.
- Wells, L. (2013). The o-mannosylation pathway: glycosyltransferases and proteins implicated in congenital muscular dystrophy. J. Biol. Chem., 288, 6930-5.
- Merlini, L., Brunner, HG., van Bokhoven, H., Chitayat, D., van Beusekom, E., Celli, J. et al. (2002). Mutations in the O-mannosyltransferase gene POMT1 give rise to the severe neuronal migration disorder Walker-Warburg syndrome. *Am. J. Hum. Genet.*, *71*, 1033-43.
- Gualandi, F., D'Amico, A., Petrini, S., Bertini, E. (2011). Congenital muscular dystrophies: a brief review. Semin Pediatr Neurol, 18, 277-88.

# Editions

2014-07-25	Authored, Edited	Jassal, B.
2015-12-18	Reviewed	Joshi, HJ., Hansen, L.