

GLI3R binds the GLI2 promoter

Gillespie, ME., Liu, Y C., Rothfels, K.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](#). For more information see our [license](#).

03/05/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 88


This document contains 1 reaction ([see Table of Contents](#))

GLI3R binds the GLI2 promoter [↗](#)

Stable identifier: R-HSA-5617410

Type: binding

Compartments: nucleoplasm



GLI3R is a DNA-binding transcriptional repressor that recognizes consensus GLI sites 5'-GACCACCC-3' in the promoters of target genes (Kinzler and Vogelstein, 1990). DNA-binding is mediated through 5 C2H2 Kruppel zinc fingers in the N-terminal region of the protein, which remains intact after proteasome-mediated processing (reviewed in Hui and Angers, 2011). In the absence of Hh signaling, GLI3R has been shown to bind to the promoters of the GLI1 and GLI2 genes as assessed by ChIP and to repress gene expression (Hu et al, 2006).

Literature references

Mo, R., Hui, CC., Hu, MC., Bhella, S., Chuang, PT., Rosenblum, ND. et al. (2006). GLI3-dependent transcriptional repression of Gli1, Gli2 and kidney patterning genes disrupts renal morphogenesis. *Development*, 133, 569-78. [↗](#)

Hui, CC., Angers, S. (2011). Gli proteins in development and disease. *Annu. Rev. Cell Dev. Biol.*, 27, 513-37. [↗](#)

Kinzler, KW., Vogelstein, B. (1990). The GLI gene encodes a nuclear protein which binds specific sequences in the human genome. *Mol. Cell. Biol.*, 10, 634-42. [↗](#)

Editions

2014-07-23	Authored	Rothfels, K.
2014-07-25	Edited	Gillespie, ME.
2014-08-01	Reviewed	Liu, Y C.