

TP53 stimulates expression of SESN1,2,3 genes

Hwang, PM., Inga, A., Kang, JG., Orlic-Milacic, M., Wang, PY., Zaccara, S.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](#). For more information see our [license](#).

03/05/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 88


This document contains 1 reaction ([see Table of Contents](#))

TP53 stimulates expression of SESN1,2,3 genes ↗

Stable identifier: R-HSA-5629189

Type: omitted

Compartments: nucleoplasm



Sestrins (SESN) are a small family of stress-sensitive gene that are conserved across several species. Mammals express three different SESN family members characterized as SESN1-3. Sestrin genes, SESN1, SESN2 and SESN3, are upregulated in response to TP53-mediated transcriptional regulation. SESN1 and SESN2 were classified as members of the growth arrest and DNA damage (GADD) gene family that can regulate cell growth and viability under different cellular pressures. In particular, p53 negatively modulates the mTOR pathway via SESN1 and SESN2 upregulation (Feng 2010). SESN3 was identified shortly after SESN2 through in silico analysis and was found to be a target of the forkhead transcription factor (FOXO) family. A specific TP53 binding site on the human SESN3 promoter has not been identified yet, but was found in the rat ortholog (Velasco-Miguel et al. 1999, Budanov et al. 2002, Brynczka et al. 2007).

Literature references

- Gelbert, L., Laidlaw, J., Talbott, R., Seizinger, B., Buckbinder, L., Jean, P. et al. (1999). PA26, a novel target of the p53 tumor suppressor and member of the GADD family of DNA damage and growth arrest inducible genes. *Oncogene*, 18, 127-37. ↗
- Feng, Z. (2010). p53 regulation of the IGF-1/AKT/mTOR pathways and the endosomal compartment. *Cold Spring Harb Perspect Biol*, 2, a001057. ↗
- Labhart, P., Brynczka, C., Merrick, BA. (2007). NGF-mediated transcriptional targets of p53 in PC12 neuronal differentiation. *BMC Genomics*, 8, 139. ↗
- Feinstein, E., Gudkov, AV., Chajut, A., Kamer, I., Kalinski, H., Fishman, A. et al. (2002). Identification of a novel stress-responsive gene Hi95 involved in regulation of cell viability. *Oncogene*, 21, 6017-31. ↗

Editions

2014-12-23	Authored, Edited	Orlic-Milacic, M.
2014-12-30	Reviewed	Hwang, PM., Kang, JG., Wang, PY.
2016-02-04	Reviewed	Inga, A., Zaccara, S.