

SORD oxidizes D-sorbitol to Fru

Cameselle, J.C., D'Eustachio, P., Hill, D.P., Jassal, B., Ribeiro, J.M.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](#). For more information see our [license](#).

02/05/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 88


This document contains 1 reaction ([see Table of Contents](#))

SORD oxidizes D-sorbitol to Fru [↗](#)

Stable identifier: R-HSA-5652195

Type: transition

Compartments: cytosol



Cytosolic SORD (sorbitol dehydrogenase) catalyzes the reaction of D-sorbitol and NAD⁺ to form fructose (Fru) and NADH + H⁺. This reaction was first described by Hers (1960) in sheep seminal vesicles; the human enzyme was identified by O'Brien et al. (1983). The active enzyme is a tetramer with four associated Zn²⁺ ions (Pauly et al. 2003) whose amino-terminal methionine residue has been removed (Karlsson et al. 1989).

Literature references

- Adams, PD., Cunningham, D., Kamath, A., Mcguire, D., Chrnyk, B., Madura, R. et al. (2003). X-ray crystallographic and kinetic studies of human sorbitol dehydrogenase. *Structure*, 11, 1071-85. [↗](#)
- Edwards, MR., Schofield, PJ., O'Brien, MM. (1983). Polyol-pathway enzymes of human brain. Partial purification and properties of sorbitol dehydrogenase. *Biochem. J.*, 211, 81-90. [↗](#)
- Höög, JO., Karlsson, C., Jörnvall, H., Maret, W., Auld, DS. (1989). Variability within mammalian sorbitol dehydrogenases. The primary structure of the human liver enzyme. *Eur. J. Biochem.*, 186, 543-50. [↗](#)
- Hers, HG. (1960). [The mechanism of the formation of seminal fructose and fetal fructose]. *Biochim. Biophys. Acta*, 37, 127-38. [↗](#)

Editions

2014-11-29	Authored, Edited	D'Eustachio, P.
2015-01-29	Reviewed	Jassal, B.
2015-08-28	Authored	Ribeiro, JM., Cameselle, JC.
2023-05-16	Reviewed	Hill, DP.