

FURIN cleaves 7K-BACE1 to 7K-BACE1(46-501)

D'Eustachio, P., Jassal, B., Perry, G.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://reactome.org/licenses/).

01/05/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 88


This document contains 1 reaction ([see Table of Contents](#))

FURIN cleaves 7K-BACE1 to 7K-BACE1(46-501) ↗

Stable identifier: R-HSA-5693081

Type: omitted

Compartments: Golgi-associated vesicle lumen, Golgi membrane



FURIN is the most likely endopeptidase that cleaves the BACE propeptide domain (BACE1(22-45)) to form the mature enzyme (7K-BACE1(46-501)). Although the pro-enzyme possesses proteolytic activity, this activity is approximately doubled following removal of the prodomain (Bennett et al. 2000).

Literature references

Citron, M., Haniu, M., Denis, P., Teplow, DB., Vassar, R., Louis, JC. et al. (2000). A furin-like convertase mediates propeptide cleavage of BACE, the Alzheimer's beta -secretase. *J. Biol. Chem.*, 275, 37712-7. ↗

Editions

2015-05-13	Authored, Edited	Jassal, B.
2015-06-26	Reviewed	D'Eustachio, P.
2015-11-09	Reviewed	Perry, G.