

Mitochondrial RNase P (mtRNase P) cleaves the 5' ends of pre-tRNAs and ELAC2 (RNase Z) cleaves the 3' ends of pre tRNAs in the H strand transcript

Jarrous, N., Levinger, L., May, B., Motorin, Y., Powell, CA., Suzuki, T.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of <u>Creative Commons Attribution 4.0 International (CC BY 4.0)</u>
<u>License</u>. For more information see our <u>license</u>.

19/05/2024

https://reactome.org

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142.
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467.
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res, 46*, D649-D655.
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph data-base: Efficient access to complex pathway data. *PLoS computational biology, 14*, e1005968.

Reactome database release: 88

This document contains 1 reaction (see Table of Contents)


https://reactome.org Page 2

Mitochondrial RNase P (mtRNase P) cleaves the 5' ends of pre-tRNAs and ELAC2 (RNase Z) cleaves the 3' ends of pre-tRNAs in the H strand transcript **↗**

Stable identifier: R-HSA-6785722

Type: omitted

Compartments: mitochondrial matrix



RNase P, ELAC2, and additional unknown nucleases cleave H strand transcripts to release the various tRNAs, rRNAs, and mRNAs contained in the long polycistronic transcripts.

Mitochondrial RNase P, comprising 3 protein subunits and no RNA moiety (Holzmann et al. 2008), endonucleolytically cleaves polycistronic mitochondrial transcripts at the 5' ends of the tRNA sequences (Sanchez et al. 2011, Howard et al. 2012, Vilardo et al. 2012, Li et al. 2015, Reinhard et al. 2015, Vilardo and Rossmanith 2015). A subcomplex of RNase P also functions as a tRNA methyltransferase and the SDR5C1 subunit is an amino acid and fatty acid dehydrogenase. Mutations in the SDR5C1 subunit of RNase P cause HSD10 disease, which is characterized by progressive neurodegeneration and cardiomyopathy (Vilardo and Rossmanith 2015)

ELAC2 cleaves polycistronic mitochondrial transcripts at the 3' ends of the tRNA sequences (Brzezniak et al. 2011, Sanchez et al. 2011). Different isoforms of ELAC2 are present in the nucleus and mitochondria (Rossmanith 2011). Mutations in ELAC2 cause cardiac hypertrophy (Haack et al. 2013) and disorders of oxidative phosphorylation (reviewed in Van Haute et al. 2015).

Unknown nucleases also cleave the H strand transcript at sites 5' to MT-CO3, 5' to MT-CO1, and 5' to MT-CYB (reviewed in Van Haute et al. 2015).

Literature references

Reinhard, L., Sridhara, S., Hallberg, BM. (2015). Structure of the nuclease subunit of human mitochondrial RNase P. *Nucleic Acids Res.*, 43, 5664-72.

Zhou, W., Li, F., Liu, X., Yang, X., Shen, Y. (2015). Auto-inhibitory Mechanism of the Human Mitochondrial RNase P Protein Complex. *Sci Rep*, 5, 9878.

Koutmos, M., Lim, WH., Fierke, CA., Howard, MJ. (2012). Mitochondrial ribonuclease P structure provides insight into the evolution of catalytic strategies for precursor-tRNA 5' processing. *Proc. Natl. Acad. Sci. U.S.A., 109*, 16149-54.

Bennett, KL., Löffler, E., Rossmanith, W., Gerner, C., Holzmann, J., Frank, P. (2008). RNase P without RNA: identification and functional reconstitution of the human mitochondrial tRNA processing enzyme. *Cell*, 135, 462-74.

Serjanov, D., Levinger, L. (2012). Pathogenesis-related mutations in the T-loops of human mitochondrial tRNAs affect 3' end processing and tRNA structure. RNA Biol, 9, 283-91.

https://reactome.org

Editions

2015-06-29	Authored, Edited	May, B.
2015-08-11	Reviewed	Levinger, L.
2015-08-25	Reviewed	Motorin, Y.
2015-10-24	Reviewed	Jarrous, N.
2015-11-07	Reviewed	Powell, CA.
2015-11-18	Reviewed	Suzuki, T.