

# DNA nucleases bind monoubiquitinated ID2 complex

Fugger, K., Orlic-Milacic, M.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://reactome.org/licenses).

14/05/2024

## Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

## Literature references

Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)

Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)

Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)

Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 88


This document contains 1 reaction ([see Table of Contents](#))

## DNA nucleases bind monoubiquitinated ID2 complex ↗

**Stable identifier:** R-HSA-6785732

**Type:** binding

**Compartments:** nucleoplasm



Several DNA nucleases bind to interstrand crosslinks (ICL-DNA) and participate in ICL "unhooking". The ubiquitin-binding zinc finger (UBZ) domain of the DNA nuclease FAN1 binds to monoubiquitinated FANCD2, enabling the recruitment of FAN1 to the ICL-DNA repair site (Liu et al. 2010, MacKay et al. 2010, Smogorzewska et al. 2010, Kratz et al. 2010). Once recruited to ICL-DNA, FAN1 forms head-to-tail homodimers. Homodimerization is important for the endonucleolytic activity of FAN1 (Zhao et al. 2014). SLX4 (FANCP) serves as a docking platform for recruitment of SLX1A, MUS81 and EME1 or EME2, resulting in formation of the SLX1A:SLX4:MUS81:EME1 (or SLX1A:SLX4:MUS81:EME2) endonucleolytic complex (Fekairi et al. 2009, Wyatt et al. 2013). SLX4 can also bind the endonucleolytic complex composed of ERCC1 and ERCC4 (XPF) (Fekairi et al. 2009). SLX4 is recruited to ICL-DNA through interaction of the UBZ domain of SLX4 with monoubiquitinated FANCD2 (Yamamoto et al. 2011). Targeted deletion of the UBZ domain of SLX4 confers sensitivity to ICL-inducing agents, but the UBZ domain seems to be dispensable for the role of SLX4 in homologous recombination repair (Yamamoto et al. 2011).

DNA exonucleases DCLRE1A (SNM1A) and DCLRE1B (SNM1B) likely function redundantly in ICL repair. Similar to FAN1, they are able to digest the DNA past the ICL, thereby unhooking one of the DNA strands (Wang et al. 2011, Sengerova et al. 2012). Monoubiquitination of the PCNA subunit of the stalled replicative polymerase complex by RAD18 may provide the docking site for DCLRE1A (or DCLRE1B) (Yang et al. 2010). In addition, PIAS1 may facilitate loading of DCLRE1A (or DCLRE1B) to ICL sites (Ishiai et al. 2004).

## Literature references

- Matsushita, N., Yamazoe, M., Takata, M., Buerstedde, JM., Takeda, S., Agematsu, K. et al. (2004). DNA cross-link repair protein SNM1A interacts with PIAS1 in nuclear focus formation. *Mol Cell Biol*, 24, 10733-41. ↗
- West, SC., Hofmann, K., Déclais, AC., Gartner, A., Deans, AJ., Lilley, DM. et al. (2010). Identification of KIAA1018/FAN1, a DNA repair nuclease recruited to DNA damage by monoubiquitinated FANCD2. *Cell*, 142, 65-76. ↗
- Longerich, S., Zhao, Q., Xue, X., Sung, P., Xiong, Y. (2014). Structural insights into 5' flap DNA unwinding and incision by the human FAN1 dimer. *Nat Commun*, 5, 5726. ↗
- Hartley, J., McHugh, PJ., Lee, SY., Gileadi, O., Abu, M., Sengerová, B. et al. (2012). Characterization of the human SNM1A and SNM1B/Apollo DNA repair exonucleases. *J. Biol. Chem.*, 287, 26254-67. ↗

West, SC., Matos, J., Sarbajna, S., Wyatt, HD. (2013). Coordinated actions of SLX1-SLX4 and MUS81-EME1 for Holliday junction resolution in human cells. *Mol. Cell*, 52, 234-47. [↗](#)

**Editions**

2015-06-29	Authored, Edited	Orlic-Milacic, M.
2015-08-20	Reviewed	Fugger, K.