

HOXB2 chromatin is activated

Blasi, F., May, B., Rezsöházy, R.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://reactome.org/licenses/).

02/05/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 88

This document contains 1 reaction ([see Table of Contents](#))


HOXB2 chromatin is activated ↗

Stable identifier: R-HSA-6810159

Type: omitted

Compartments: nucleoplasm

Inferred from: [Hoxb2 chromatin is activated \(Mus musculus\)](#)



During activation of HOXB1 by retinoic acid in human embryonal carcinoma cells, methylation at lysine-27 of histone H3 (H3K27me3) is lost and methylation at lysine-4 (H3K4me3) is gained (Lan et al. 2007, Lee et al. 2007). The histone demethylase KDM6A (UTX) binds HOXB2 chromatin and may demethylate H3K27me3 (Lee et al. 2007). Other factors may also participate in demethylation. Loss of H3K27me3 is associated with loss of polycomb repressive complex 2 (PRC2) (Lan et al. 2007, Lee et al. 2007). KDM6A forms complexes with the histone methyltransferases KMT2C,D (MLL2,3) which may participate in methylating H3K4 (Lee et al. 2007). The activation of HOXB1 chromatin may be produced by euchromatin spreading from distant 3' retinoic acid response elements.

Literature references

- Rinn, JL., Wang, JK., Lan, F., Shi, Y., Issaeva, I., Chang, HY. et al. (2007). A histone H3 lysine 27 demethylase regulates animal posterior development. *Nature*, 449, 689-94. ↗
- Di Croce, L., Trojer, P., Reinberg, D., Lee, MG., Villa, R., Yan, KP. et al. (2007). Demethylation of H3K27 regulates polycomb recruitment and H2A ubiquitination. *Science*, 318, 447-50. ↗

Editions

2014-08-08	Authored	May, B.
2015-05-13	Reviewed	Blasi, F.
2015-11-15	Edited	May, B.
2015-11-15	Authored	Rezsohazy, R.