

Synaptic adhesion-like molecules



Garapati, P V., Kim, E., Petralia, RS., Seabold, GK.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of <u>Creative Commons Attribution 4.0 International (CC BY 4.0)</u> <u>License</u>. For more information see our <u>license</u>.

This is just an excerpt of a full-length report for this pathway. To access the complete report, please download it at the <u>Reactome Textbook</u>.

06/05/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics, 18*, 142. 7
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. A
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res, 46*, D649-D655. ↗
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, *14*, e1005968. *オ*

This document contains 1 pathway and 8 reactions (see Table of Contents)

Synaptic adhesion-like molecules 7

Stable identifier: R-HSA-8849932



Recruitment of receptors and ion channels to the postsynaptic membrane is the last step in synapse formation. Many of these proteins interact directly or indirectly with postsynaptic density-95 (PSD95)/Discs large/zona occludens-1 (PDZ) proteins, thus linking them to the postsynaptic scaffold and providing a mechanism for both retaining the protein at the synapse and keeping its proximity to signaling molecules known to associate with PDZ proteins (Wang et al. 2006, Morimura et al. 2006, Ko et al. 2006, Nourry et al. 2003, Kim & Sheng 2004, Montgomery et al. 2004, Sheng and Kim 2011). The synaptic adhesion-like molecules (SALM) family belongs to the superfamily of leucine-rich repeat (LRR)-containing adhesion molecules, alternatively referred to as LRFN (leucine-rich repeat and fibronectin III domain-containing) is synapse adhesion molecule linked to NMDA and AMPA receptors. It includes five known members (SALMs 1-5 or LRFN1-5), which have been implicated in the regulation of neurite outgrowth and branching, and synapse formation and maturation. SALM proteins are distributed to both dendrites and axons in neurons (Ko et al. 2006, Wang et al. 2006, Sebold et al. 2012). The family members, SALM1-SALM5, have a single transmembrane (TM) domain and contain extracellular leucine-rich repeats, an Ig C2 type domain, a fibronectin type III domain, and an intracellular postsynaptic density-95 (PSD-95)/Discs large/zona occludens-1 (PDZ) binding domain, which is present on all members except SALM4 and SALM5 (Ko et al. 2006, Wang et al. 2006, Morimura et al. 2006).

Literature references

- Kim, E., Mah, W., Nam, J. (2011). The SALM/Lrfn family of leucine-rich repeat-containing cell adhesion molecules. Semin. Cell Dev. Biol., 22, 492-8. ↗
- Kaang, BK., Kim, H., Kim, K., Chung, HS., Kim, E., Ko, J. et al. (2006). SALM synaptic cell adhesion-like molecules regulate the differentiation of excitatory synapses. *Neuron*, *50*, 233-45. *¬*

Kim, E., Sheng, M. (2011). The postsynaptic organization of synapses. Cold Spring Harb Perspect Biol, 3. 🛪

2015-12-18	Authored, Edited	Garapati, P V.
2016-02-04	Reviewed	Kim, E.
2016-05-17	Reviewed	Petralia, RS., Seabold, GK.

PSD-95 binds NMDA receptor ↗

Location: Synaptic adhesion-like molecules

Stable identifier: R-HSA-8849878

Type: binding

Compartments: plasma membrane



Many membrane proteins that contain a PDZ-binding domain require an interaction with a PDZ protein for correct targeting to the cell surface (Zheng et al. 2011). N-methyl-D-aspartate (NMDA) receptor interact with postsynaptic density protein (PSD-95) and the PSD-95 family might serve to anchor NMDA receptors to the submembrane cytoskeleton and aid in the assembly of signal transduction complexes at postsynaptic sites (Niethammer et al. 1996, Kornau et al. 1995, Sans et al. 2000).

Literature references

Sheng, M., Kim, E., Niethammer, M. (1996). Interaction between the C terminus of NMDA receptor subunits and multiple members of the PSD-95 family of membrane-associated guanylate kinases. J. Neurosci., 16, 2157-63.

Schenker, LT., Kornau, HC., Seeburg, PH., Kennedy, MB. (1995). Domain interaction between NMDA receptor subunits and the postsynaptic density protein PSD-95. *Science*, 269, 1737-40.

2015-12-18	Authored, Edited	Garapati, P V.
2016-02-04	Reviewed	Kim, E.
2016-05-17	Reviewed	Petralia, RS., Seabold, GK.

SALM1 binds NMDA receptor 7

Location: Synaptic adhesion-like molecules

Stable identifier: R-HSA-8849906

Type: binding

Compartments: plasma membrane



SALM1 interacts with and recruits NMDA receptors to early synapses. NMDA receptors are involved in the development of excitatory synapse by interacting with PDZ proteins of the PSD-95 family through its NR2 subunits. SALM1 can directly interact with the extracellular domain of the NR1 subunit of NMDA receptor or indirectly by binding to PSD-95, which may recruit NMDA receptor via the NR2 subunits of NMDA receptors (Wang et al. 2006, Niethammer et al. 1996, Kornau et al. 1995). SALM1 also often localizes with the NR1 subunit of NMDA receptors in hippocampal and cerebellar neurons (Thevenon et al., 2015).

Literature references

- Schenker, LT., Kornau, HC., Seeburg, PH., Kennedy, MB. (1995). Domain interaction between NMDA receptor subunits and the postsynaptic density protein PSD-95. *Science*, 269, 1737-40.
- Wang, YX., Wang, CY., Wenthold, RJ., Petralia, RS., Seabold, GK., Chang, K. (2006). A novel family of adhesion-like molecules that interacts with the NMDA receptor. *J. Neurosci.*, *26*, 2174-83.

2015-12-18	Authored, Edited	Garapati, P V.
2016-02-04	Reviewed	Kim, E.
2016-05-17	Reviewed	Petralia, RS., Seabold, GK.

SALMs 1-3 bind to PSD-95 family members 7

Location: Synaptic adhesion-like molecules

Stable identifier: R-HSA-8849891

Type: binding

Compartments: plasma membrane



SALMs 1-3 interact with the PDZ domain containing proteins PSD 95 (DLG4) and synapse associated protein 97 (SAP97 or DLG1) and SAP102 (DLG3), based on yeast 2-hybrid assays (Wang et al. 2006, Ko et al. 2006) and coimmunoprecipitations from detergent solubilized brain (Wang et al. 2006, Ko et al. 2006, Mah et al. 2010) and transiently transfected mammalian cells (Morimura et al., 2006, Wang et al. 2006). PDZ proteins play a central role in organizing functionally diverse membrane proteins at the synapse (Wang et al. 2006, Zheng et al. 2011). PSD-95 family members are abundant postsynaptic scaffolding proteins at excitatory synapses. SALM1 (Wang et al. 2006), SALM2 (Ko et al. 2006), SALM3 and SALM5 (Mah et al. 2010) proteins are enriched in synaptic fractions. SALM5 forms a weak complex with PSD-95, an abundant postsynaptic scaffolding protein at excitatory synapse most likely through indirect interactions (Mah et al. 2010). SALM3 and SALM5, but not other SALMs, induce presynaptic differentiation in contacting axons (Mah et al. 2010).

Literature references

Wang, YX., Wang, CY., Wenthold, RJ., Petralia, RS., Seabold, GK., Chang, K. (2006). A novel family of adhesion-like molecules that interacts with the NMDA receptor. J. Neurosci., 26, 2174-83.

2015-12-18	Authored, Edited	Garapati, P V.
2016-02-04	Reviewed	Kim, E.
2016-05-17	Reviewed	Petralia, RS., Seabold, GK.

SALMs1-3 bind each other in cis interactions *▼*

Location: Synaptic adhesion-like molecules

Stable identifier: R-HSA-8849900

Type: binding

Compartments: plasma membrane



SALM1, SALM2, and SALM3 form homo- and heteromeric complexes in a cis manner. These SALMs 1-3 coimmunoprecipitate with each other in extracts from brain. Whether the homo and heteromeric complexes formed between SALMs 1-3 contribute to synapse formation or neurite outgrowth remains to be determined (Seabold et al. 2008).

Literature references

Wang, YX., Wang, CY., Wang, PY., Wenthold, RJ., Petralia, RS., Chang, K. et al. (2008). The SALM family of adhesionlike molecules forms heteromeric and homomeric complexes. J. Biol. Chem., 283, 8395-405.

2015-12-18	Authored, Edited	Garapati, P V.
2016-02-04	Reviewed	Kim, E.
2016-05-17	Reviewed	Petralia, RS., Seabold, GK.

SALM2 associate with AMPA and NMDA receptors 7

Location: Synaptic adhesion-like molecules

Stable identifier: R-HSA-8849881

Type: binding

Compartments: plasma membrane



SALM2 coimmunoprecipitates with NMDAR and AMPAR subunits isolated from detergent-solubilized brain (Ko et al. 2006). SALM2 co localizes with both pre- and post-synaptic proteins at excitatory synapses in mature neurons. SALM2 associates with AMPA receptors and, to a lesser extent, with NMDA receptors in hippocampal cultures. SALM2 appears to promote the maturation, but not the formation, of excitatory synapses, in line with the fact that synaptic localization of AMPA receptors occurs at late stages of excitatory synaptic development (Ko et al. 2006, Nam et al. 2011).

Literature references

Kaang, BK., Kim, H., Kim, K., Chung, HS., Kim, E., Ko, J. et al. (2006). SALM synaptic cell adhesion-like molecules regulate the differentiation of excitatory synapses. *Neuron*, *50*, 233-45.

2015-12-18	Authored, Edited	Garapati, P V.
2016-02-04	Reviewed	Kim, E.
2016-05-17	Reviewed	Petralia, RS., Seabold, GK.

SALMs 1-4 bind reticulon 3 ↗

Location: Synaptic adhesion-like molecules

Stable identifier: R-HSA-8849882

Type: binding

Compartments: plasma membrane



In the brain, reticulon 3 (RTN3) tightly associates with SALM2 and SALM3 to form a complex, and interacts relatively weakly with SALM1 and SALM4. This interaction is mediated by the LRR domain in SALMs (Chang et al. 2010).

Literature references

Wang, CY., Wenthold, RJ., Seabold, GK., Chang, K. (2010). Reticulon 3 is an interacting partner of the SALM family of adhesion molecules. J. Neurosci. Res., 88, 266-74. 🛪

2015-12-18	Authored, Edited	Garapati, P V.
2016-02-04	Reviewed	Kim, E.
2016-05-17	Reviewed	Petralia, RS., Seabold, GK.

SALM4 binds flotillin-1 ↗

Location: Synaptic adhesion-like molecules

Stable identifier: R-HSA-8849908

Type: binding

Compartments: plasma membrane



SALM4 induces neurite branching by binding to flotillin-1, a lipid raft-associated protein that interacts with NMDA receptors (Swanwick et al. 2009, 2010).

Literature references

Shapiro, ME., Wenthold, RJ., Swanwick, CC., Vicini, S. (2010). Flotillin-1 mediates neurite branching induced by synaptic adhesion-like molecule 4 in hippocampal neurons. *Mol. Cell. Neurosci.*, 45, 213-25.

2015-12-18	Authored, Edited	Garapati, P V.
2016-02-04	Reviewed	Kim, E.
2016-05-17	Reviewed	Petralia, RS., Seabold, GK.

SALM3 binds LAR-RPTP ↗

Location: Synaptic adhesion-like molecules

Stable identifier: R-HSA-8855648

Type: binding

Compartments: plasma membrane



SALM3 interacts with LAR family receptor protein tyrosine phosphatases (LAR-RPTPs) in a transynaptic manner that is dependent upon a splice insert in LAR-RPTPs. This interaction regulates SALM3 dependent presynaptic differentiation (Li et al. 2015). SALM3 knockout mice have fewer excitatory synapsese, but normal plasticity in the hippocampus. However, they are hypoactive, suggesting SALM3 influences locomotion behavior (Li et al. 2015).

Literature references

Choi, SY., Zhang, P., Craig, AM., Kim, Y., Roh, JD., Bae, YC. et al. (2015). Splicing-Dependent Trans-synaptic SALM3-LAR-RPTP Interactions Regulate Excitatory Synapse Development and Locomotion. *Cell Rep, 12*, 1618-30. ↗

2015-12-18	Authored, Edited	Garapati, P V.
2016-02-04	Reviewed	Kim, E.
2016-05-17	Reviewed	Petralia, RS., Seabold, GK.

Table of Contents

Introduction	1
📱 Synaptic adhesion-like molecules	2
▶ PSD-95 binds NMDA receptor	3
➢ SALM1 binds NMDA receptor	4
➢ SALMs 1-3 bind to PSD-95 family members	5
> SALMs1-3 bind each other in cis interactions	6
> SALM2 associate with AMPA and NMDA receptors	7
⇒ SALMs 1-4 bind reticulon 3	8
⇒ SALM4 binds flotillin-1	9
⇒ SALM3 binds LAR-RPTP	10
Table of Contents	11