

CBFB binds RUNX3

Chuang, L.S., Ito, Y., Orlic-Milacic, M.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://reactome.org/licenses/).

03/05/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 88


This document contains 1 reaction ([see Table of Contents](#))

CBFB binds RUNX3 [↗](#)

Stable identifier: R-HSA-8865454

Type: binding

Compartments: nucleoplasm



CBFB binds the transcription factor RUNX3, interacting with its Runt domain, which is highly similar to Runt domains of RUNX1 and RUNX2 (Kim et al. 2013). RUNX3 is implicated in neurogenesis, thymopoiesis and stomach development (Inoue et al. 2002, Levanon et al. 2002, Levanon et al. 2003). RUNX3 and CBFB are frequently downregulated in gastric cancer (Sakakura et al. 2005).

Literature references

Hartley, PD., Mann, S., Gross, JD., Krogan, NJ., Kim, DY., Crosby, DC. et al. (2013). CBF β stabilizes HIV Vif to counteract APOBEC3 at the expense of RUNX1 target gene expression. *Mol. Cell*, 49, 632-44. [↗](#)

Editions

2016-12-13	Authored	Orlic-Milacic, M.
2017-01-31	Reviewed	Ito, Y., Chuang, LS.
2017-01-31	Edited	Orlic-Milacic, M.