

Cleaved fibrinogen products bind

TLR4:LY96

Granucci, F., Shamovsky, V., Zanoni, I.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of <u>Creative Commons Attribution 4.0 International (CC BY 4.0)</u> <u>License</u>. For more information see our <u>license</u>.

17/05/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics, 18,* 142. 7
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. A
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res, 46*, D649-D655. ↗
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, *14*, e1005968. *オ*

This document contains 1 reaction (see Table of Contents)

Cleaved fibrinogen products bind TLR4:LY96 7

Stable identifier: R-HSA-8870678

Type: binding

Compartments: plasma membrane



Fibrinogen, in addition to its role in coagulation, is also an acute phase protein of inflammation which can induce a cytokine production acting as an endogenous ligand for toll-like receptor 4 (TLR4) expressed on cells including macrophages and airway epithelial cells (Millien VO et al., 2013; Kuhns DB et al., 2007; Smiley ST et al., 2001). In human macrophages fibrinogen stimulated interleukin IL6 expression and extracellular signal-related kinase (ERK) phosphorylation ((Hodgkinson CP et al., 2008). In human embryonic kidney 293 (HEK293)-CD14-MD2 cells expressing TLR4, fibrinogen induced robust phosphorylation of ERK1, p38alpha and JNK and activated transcription factors NFkappaB, Elk1 and AP1 (Hodgkinson CP et al., 2008). Moreover, proteinases, such as thrombin, can cleave fibrinogen. In mice, exposure to endogenous or exogenous proteinases lead to hyperactivation of an antifungal pathway and lead to allergic airway inflammation through activation of TLR4-dependent signaling pathway by fibrinogen cleaved products (Millien VO et al., 2013)

Literature references

Hancock, WW., Smiley, ST., King, JA. (2001). Fibrinogen stimulates macrophage chemokine secretion through tolllike receptor 4. *J Immunol*, *167*, 2887-94. *对*

Editions

2016-05-10	Authored, Edited	Shamovsky, V.
2016-05-12	Reviewed	Zanoni, I., Granucci, F.