

E2F6 binds TFDP1

Herlihy, A., Orlic-Milacic, M.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://reactome.org/licenses/).

29/04/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 88


This document contains 1 reaction ([see Table of Contents](#))

E2F6 binds TFDP1 [↗](#)

Stable identifier: R-HSA-8953111

Type: binding

Compartments: nucleoplasm



E2F6, an E2F family member without the pocket protein-binding domain and transactivation domain (Gaubatz et al. 1998), forms a heterodimer with TFDP1 (DP-1) (Cartwright et al. 1998, Trimarchi et al. 1998, Ogawa et al. 2002).

Literature references

Livingston, DM., Gaubatz, S., Ishiguro, K., Ogawa, H., Nakatani, Y. (2002). A complex with chromatin modifiers that occupies E2F- and Myc-responsive genes in G0 cells. *Science*, 296, 1132-6. [↗](#)

Holm, K., Wagener, C., Helin, K., Müller, H., Cartwright, P. (1998). E2F-6: a novel member of the E2F family is an inhibitor of E2F-dependent transcription. *Oncogene*, 17, 611-23. [↗](#)

Verona, R., Andon, N., Moberg, K., Lees, JA., Fairchild, B., Trimarchi, JM. (1998). E2F-6, a member of the E2F family that can behave as a transcriptional repressor. *Proc. Natl. Acad. Sci. U.S.A.*, 95, 2850-5. [↗](#)

Editions

2017-01-05	Authored	Orlic-Milacic, M.
2017-05-12	Reviewed	Herlihy, A.
2017-05-19	Edited	Orlic-Milacic, M.