

COMMDs displace CAND1 from CRL4 E3 ubiquitin ligase complex

Pick, E., Rothfels, K.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://reactome.org/licenses/).

04/05/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 88


This document contains 1 reaction ([see Table of Contents](#))

COMMDs displace CAND1 from CRL4 E3 ubiquitin ligase complex ↗

Stable identifier: R-HSA-8955285

Type: transition

Compartments: nucleoplasm



COMMD1 is a member of a family of 10 copper metabolism MURR1 domain-containing proteins that have pleiotropic roles in copper metabolism, NF kappa beta-mediated transcription, the hypoxic response and electrolyte transport (Burstein et al, 2005; reviewed in Maine and Burstein, 2007). COMMD proteins have differential tissue and expression levels, but appear to have partially overlapping function and form homo- and heterodimers through the shared COMM domain (Burstein et al, 2005). COMMD1 and other family members interact with the cullin subunit of CRL E3 ubiquitin ligase complexes, as well as with CCDC22, a protein implicated in X-linked intellectual disability that may regulate COMMD localization. Together, COMMD proteins and CCDC22 activate the ubiquitin ligase activity of CRL complexes by displacing the CAND1 inhibitor (Burstein et al, 2005; Maine et al, 2007; Mao et al, 2011; Starokadomskyy et al, 2013;Phillips-Krawczak et al, 2015). The specificity of interaction between various COMMD and CUL family members may serve to fine tune the regulation of CRL activation, although these details remain to be determined.

Literature references

- Maine, GN., Mao, X., Komarck, CM., Burstein, E. (2007). COMMD1 promotes the ubiquitination of NF-kappaB subunits through a cullin-containing ubiquitin ligase. *EMBO J.*, 26, 436-47. ↗
- Gluck, N., Maine, GN., Li, H., Starokadomskyy, P., Chen, B., Mao, X. et al. (2011). COMMD1 (copper metabolism MURR1 domain-containing protein 1) regulates Cullin RING ligases by preventing CAND1 (Cullin-associated Nedd8-dissociated protein 1) binding. *J. Biol. Chem.*, 286, 32355-65. ↗
- Gluck, N., Zecha, A., Maine, GN., Li, H., Wallis, M., Ropers, HH. et al. (2013). CCDC22 deficiency in humans blunts activation of proinflammatory NF-κB signaling. *J. Clin. Invest.*, 123, 2244-56. ↗
- Maine, GN., Burstein, E. (2007). COMMD proteins: COMMMing to the scene. *Cell. Mol. Life Sci.*, 64, 1997-2005. ↗
- Hoberg, JE., Maine, GN., Duckett, CS., Wilkinson, AS., Wilkinson, JC., Rumble, JM. et al. (2005). COMMD proteins, a novel family of structural and functional homologs of MURR1. *J. Biol. Chem.*, 280, 22222-32. ↗

Editions

2016-12-13	Authored, Edited	Rothfels, K.
2017-02-22	Reviewed	Pick, E.