

IL15:IL15RA:p-Y-IL2RB:p-Y-JAK1:p-Y-IL2RG:p-Y-JAK3 binds STAT3 and STAT5

Duenas, C., Patidar, M.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://reactome.org/licenses/).

03/05/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 88


This document contains 1 reaction ([see Table of Contents](#))

IL15:IL15RA:p-Y-IL2RB:p-Y-JAK1:p-Y-IL2RG:p-Y-JAK3 binds STAT3 and STAT5 [↗](#)

Stable identifier: R-HSA-8983378

Type: uncertain

Compartments: cytosol, extracellular region, plasma membrane



Signal transducer and activator of transcription 3 (STAT3), Signal transducer and activator of transcription 5A and 5B (STAT5) are believed to bind the Interleukin-15 (IL15) receptor complex, which consists of IL15, Interleukin-15 receptor alpha subunit (IL15RA, IL15R α), Interleukin-2 receptor beta subunit (IL2RB, IL2R β), which is associated with JAK1 and Interleukin receptor gamma subunit, which is associated with JAK3 (Johnston et al. 1995). This is a black-box event because STAT binding is inferred from their subsequent phosphorylation.

Literature references

Ortaldo, JR., Finbloom, DS., Shibuya, K., Kaplan, D., Giri, JD., Bacon, CM. et al. (1995). Tyrosine phosphorylation and activation of STAT5, STAT3, and Janus kinases by interleukins 2 and 15. *Proc Natl Acad Sci U S A*, 92, 8705-9. [↗](#)

Editions

2017-08-07	Authored	Duenas, C.
2017-08-07	Reviewed	Patidar, M.
2017-08-09	Edited	Duenas, C.