

sABO-B:Mn²⁺ transfers Gal to H antigen- sec to form B antigen

Jassal, B., Matsui, T.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://reactome.org/licenses/).

17/05/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 88


This document contains 1 reaction ([see Table of Contents](#))

sABO-B:Mn2+ transfers Gal to H antigen-sec to form B antigen ↗

Stable identifier: R-HSA-9034053

Type: transition

Compartments: extracellular region



As well as being a Golgi membrane resident, the histo-blood group ABO system transferase (ABO) can be proteolytically processed by an unknown protease into a soluble form, fucosylglycoprotein alpha-N-acetylgalactosaminyltransferase (sABO). A, B and AB individuals express glycosyltransferase activities that convert the H antigen to the A antigen (by addition of GalNAc), to the B antigen (by addition of Gal) or to the AB antigen (by the addition of both GalNAc and Gal). O group individuals lack such activity. Differences in four critical amino acids (176, 235, 266 and 268) alter the specificity from an A to a B glycosyltransferase (Yamamoto et al. 1990, Yamamoto & McNeill 1996, Seto et al. 1999, Alfaro et al. 2008). The soluble form of histo-blood group B transferase (sABO-B) utilises UDP-Gal to transfer galactose (Gal) to the H antigen formed via Type 1 chains to form the B antigen in secretors (B antigen-sec) (Patenaude et al. 2002, Persson et al. 2007).

Literature references

- Marken, J., White, T., Yamamoto, F., Hakomori, S., Clausen, H. (1990). Molecular genetic basis of the histo-blood group ABO system. *Nature*, 345, 229-33. ↗
- Seto, NO., Bai, Y., Borisova, SN., Persson, M., Alfaro, JA., Evans, SV. et al. (2008). ABO(H) blood group A and B glycosyltransferases recognize substrate via specific conformational changes. *J. Biol. Chem.*, 283, 10097-108. ↗
- Marcus, SL., Seto, NO., Borisova, SN., Evans, SV., Patenaude, SI., Szpacenko, A. et al. (2002). The structural basis for specificity in human ABO(H) blood group biosynthesis. *Nat. Struct. Biol.*, 9, 685-90. ↗
- Yamamoto, F., McNeill, PD. (1996). Amino acid residue at codon 268 determines both activity and nucleotide-sugar donor substrate specificity of human histo-blood group A and B transferases. In vitro mutagenesis study. *J. Biol. Chem.*, 271, 10515-20. ↗
- Evans, SV., Bundle, DR., Palcic, MM., Compston, CA., Seto, NO., Narang, SA. (1999). Donor substrate specificity of recombinant human blood group A, B and hybrid A/B glycosyltransferases expressed in Escherichia coli. *Eur. J. Biochem.*, 259, 770-5. ↗

Editions

2018-01-02	Authored, Edited	Jassal, B.
2018-12-17	Reviewed	Matsui, T.