

# Expression of SRY in testis differentiation

Imaimatsu, K., Kanai, Y., May, B.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://reactome.org/licenses/).

05/05/2024

## Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

## Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 88

This document contains 1 reaction ([see Table of Contents](#))


## Expression of SRY in testis differentiation [↗](#)

**Stable identifier:** R-HSA-9692241

**Type:** uncertain

**Compartments:** nucleoplasm

**Inferred from:** [Expression of Sry in testis differentiation \(Mus musculus\)](#)



In the gonadal ridge (genital ridge), the SRY gene is transcribed to yield mRNA and the mRNA is translated to yield SRY protein (De Santa Barbara et al. 2001, Hossain and Saunders 2001, Miyamoto et al. 2008, also inferred from mouse homologs). Transcription of SRY is activated by WT1 (Hossain and Saunders 2001, Miyamoto et al. 2008), GATA4:ZFPM2 (also called GATA4:FOG2) (Miyamoto et al. 2008), and NR5A1 (also called SF1) (De Santa Barbara et al. 2001).

### Literature references

Taniguchi, H., Viger, RS., Miyamoto, Y., Silversides, DW., Hamel, F. (2008). A GATA4/WT1 cooperation regulates transcription of genes required for mammalian sex determination and differentiation. *BMC Mol. Biol.*, 9, 44. [↗](#)

Moniot, B., De Santa Barbara, P., Berta, P., Méjean, C., Boizet-Bonhoure, B., Malclès, MH. (2001). Steroidogenic factor-1 contributes to the cyclic-adenosine monophosphate down-regulation of human SRY gene expression. *Biol. Reprod.*, 64, 775-83. [↗](#)

Hossain, A., Saunders, GF. (2001). The human sex-determining gene SRY is a direct target of WT1. *J. Biol. Chem.*, 276, 16817-23. [↗](#)

### Editions

2020-06-22

Authored, Edited

May, B.

2020-10-29

Reviewed

Kanai, Y., Imaimatsu, K.