

nsp16 acts as a cap 2'-O-methyltransferase to modify SARS-CoV-2 mRNAs

Acencio, ML., Orlic-Milacic, M., Senff-Ribeiro, A.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of <u>Creative Commons Attribution 4.0 International (CC BY 4.0)</u> <u>License</u>. For more information see our <u>license</u>.

16/05/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics, 18*, 142. 7
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. A
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res, 46*, D649-D655. ↗
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, *14*, e1005968. *オ*

This document contains 1 reaction (see Table of Contents)

nsp16 acts as a cap 2'-O-methyltransferase to modify SARS-CoV-2 mRNAs 7

Stable identifier: R-HSA-9694499

Type: transition

Compartments: cytosol, double membrane vesicle viral factory outer membrane

Diseases: COVID-19



The function of the non-structural protein nsp16 as a 2'O-methyltransferase that acts in complex with its co-activator nsp10 is conserved in SARS-CoV-2 (Viswanathan et al. 2020).

The subgenomic mRNAs of SARS-CoV-2 coronavirus are presumed to be capped at their 5' ends, based on studies of the mouse hepatitis virus (MHV) (Lai and Stohlman 1981) and the equine torovirus (van Vliet et al. 2002). 2'-O methylation of Cap-0 by SARS-CoV-2 nsp10/nsp16 has been shown, and the process is dependent on the presence of divalent metal cations (Wilamowski et al, 2021; Benoni et al, 2021; Minasov et al, 2021; Viswanathan et al, 2021). The non-structural protein 16 (nsp16) acts as a 2'O-methyltransferase that converts coronavirus cap-0 to cap-1, which was first demonstrated with nsp16 cloned from the feline coronavirus (FCV) (Decroly et al. 2008). Cap-0 represents N7-methyl guanosine connected to the 5' nucleotide through a 5' to 5' triphosphate linkage (also known as m7G cap or m7Gppp cap). Cap-1 is generated by an additional methylation on the 2'O position of the initiating nucleotide, and is also known as m7GpppNm. The non-structural protein 10 (nsp10) acts as an activator of nsp16 and is necessary for cap-1 synthesis (Bouvet et al. 2010, Decroly et al. 2011). Coronavirus RNAs with cap-1 are protected from IFIT-mediated interferon response, as IFITs recognize unmethylated 2'-O RNAs. IFITs are interferon-induced proteins with tetratricopeptide repeats that recognize unmethylated 2'-O RNAs and act to inhibit expression of virally encoded mRNAs (Menachery et al. 2014).

Literature references

- Boura, E., Krafcikova, P., Benoni, R., Baranowski, MR., Kowalska, J., Cahová, H. (2021). Substrate Specificity of SARS-CoV-2 Nsp10-Nsp16 Methyltransferase. *Viruses, 13.* 7
- Dai, N., Martínez-Sobrido, L., Chan, SH., Qi, S., Misra, A., Gupta, YK. et al. (2021). A metal ion orients SARS-CoV-2 mRNA to ensure accurate 2'-O methylation of its first nucleotide. *Nat Commun*, *12*, 3287.
- Brunzelle, JS., Minasov, G., Rosas-Lemus, M., Daczkowski, CM., Shuvalova, L., Hoover, P. et al. (2021). Mn²⁺ coordinates Cap-0-RNA to align substrates for efficient 2'-<i>O</i>-methyl transfer by SARS-CoV-2 nsp16. *Sci Signal. ¬*
- Dai, N., Martínez-Sobrido, L., Kovalskyy, D., Chan, SH., Oladunni, F., Qi, S. et al. (2020). Structural basis of RNA cap modification by SARS-CoV-2. *Nat Commun, 11*, 3718. 7

Saint, N., Minasov, G., Joachimiak, A., Rosas-Lemus, M., Shuvalova, L., Jedrzejczak, R. et al. (2021). 2'-O methylation of RNA cap in SARS-CoV-2 captured by serial crystallography. *Proc Natl Acad Sci U S A*, 118.

Editions

2020-08-12	Authored	Orlic-Milacic, M., Senff-Ribeiro, A.
2020-08-18	Edited	Orlic-Milacic, M.
2020-09-09	Reviewed	Acencio, ML.