

NeuroD1- and Pdx1-dependent synthesis of glucokinase (Gck) protein

Ferrer, J., Jensen, J., Tello-Ruiz, MK.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://reactome.org/licenses/).

29/04/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 88


This document contains 1 reaction ([see Table of Contents](#))

NeuroD1- and Pdx1-dependent synthesis of glucokinase (Gck) protein ↗

Stable identifier: R-MMU-186510

Type: omitted

Compartments: nucleoplasm, cytosol



NeuroD1 binds to control sequences of the mouse GK gene *in vitro*, and this binding is sensitive to mutagenesis in transient transfection analyses (Cissell et al. 2003; Moates et al. 2003). Moreover, NeuroD1 was shown to bind to the GK promoter *in vivo* by ChIP analysis and stimulate expression in pancreatic beta-cells (Moates et al. 2003).

Whether Pdx1 transactivates the glucokinase (GK) gene *in vivo* has been a controversial issue. Nevertheless, Pdx1 binds to control sequences of the human and mouse GK gene *in vitro*, and this binding is sensitive to mutagenesis in transient transfection analyses (Watada et al. 1996; Cissell et al. 2003). However, whereas Cissell et al. (2003) observed that Pdx1-specific antibodies against both the N and C termini of PDX1 selectively immunoprecipitate regulatory sequences of the GK gene, Chakrabarti et al. (2002) were unable to detect this association using a different Pdx1 antiserum. Disparity in results might be due to the different animal and cell line models used to study this regulation.

Literature references

- Zhao, L., Stein, R., Henderson, E., Sussel, L., Cissell, MA. (2003). Transcription factor occupancy of the insulin gene *in vivo*. Evidence for direct regulation by Nkx2.2. *J Biol Chem*, 278, 751-6. ↗
- Chakrabarti, SK., James, JC., Mirmira, RG. (2002). Quantitative assessment of gene targeting *in vitro* and *in vivo* by the pancreatic transcription factor, Pdx1. Importance of chromatin structure in directing promoter binding. *J Biol Chem*, 277, 13286-93. ↗
- Stein, R., Tsai, MJ., Cissell, MA., Nanda, S., Moates, JM. (2003). BETA2 activates transcription from the upstream glucokinase gene promoter in islet beta-cells and gut endocrine cells. *Diabetes*, 52, 403-8. ↗
- Watada, H., Yamasaki, Y., Kamada, T., Fujitani, Y., Umayahara, Y., Matsuoka, T. et al. (1996). The human glucokinase gene beta-cell-type promoter: an essential role of insulin promoter factor 1/PDX-1 in its activation in HIT-T15 cells. *Diabetes*, 45, 1478-88. ↗

Editions

2006-12-20	Edited	Tello-Ruiz, MK.
2008-05-13	Reviewed	Jensen, J.
2008-05-24	Authored	Tello-Ruiz, MK., Ferrer, J.