


Cellular responses to stress



European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of CC BY 4.0)
<u>License.</u> For more information see our License.

This is just an excerpt of a full-length report for this pathway. To access the complete report, please download it at the Reactome-Textbook.

23/04/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142.
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467.
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655.
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph data-base: Efficient access to complex pathway data. *PLoS computational biology, 14*, e1005968.


Reactome database release: 88

This document contains 10 pathways (see Table of Contents)

Cellular responses to stress ₹

Stable identifier: R-MMU-2262752

Inferred from: Cellular responses to stress (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

More details and caveats of the event inference in Reactome. For details on PANTHER see also: http://www.pantherdb.org/about.jsp


Cellular response to hypoxia >

Location: Cellular responses to stress

Stable identifier: R-MMU-1234174

Compartments: nucleoplasm, cytosol

Inferred from: Cellular response to hypoxia (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.


More details and caveats of the event inference in Reactome. For details on PANTHER see also: http://www.pantherdb.org/about.jsp

Cellular response to heat stress ↗

Location: Cellular responses to stress

Stable identifier: R-MMU-3371556

Inferred from: Cellular response to heat stress (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.


 $\underline{More\ details\ and\ cave ats\ of\ the\ event\ inference\ in\ Reactome.}\ For\ details\ on\ PANTHER\ see\ also: \\ \underline{http://www.pantherdb.org/about.jsp}$

Cellular Senescence

Location: Cellular responses to stress

Stable identifier: R-MMU-2559583

Inferred from: Cellular Senescence (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

More details and caveats of the event inference in Reactome. For details on PANTHER see also: http://www.pantherdb.org/about.jsp


HSP90 chaperone cycle for steroid hormone receptors (SHR) in the presence of ligand **→**

Location: Cellular responses to stress

Stable identifier: R-MMU-3371497

Compartments: cytosol

Inferred from: HSP90 chaperone cycle for steroid hormone receptors (SHR) in the presence of ligand (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

More details and caveats of the event inference in Reactome. For details on PANTHER see also: http://www.pantherdb.org/about.jsp


Heme signaling ₹

Location: Cellular responses to stress

Stable identifier: R-MMU-9707616

Compartments: nuclear envelope, plasma membrane, nucleoplasm, extracellular region, cytosol

Inferred from: Heme signaling (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.


More details and caveats of the event inference in Reactome. For details on PANTHER see also: http://www.pantherdb.org/about.jsp

Cellular response to chemical stress >

Location: Cellular responses to stress

Stable identifier: R-MMU-9711123

Inferred from: Cellular response to chemical stress (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

More details and caveats of the event inference in Reactome. For details on PANTHER see also: http://www.pantherdb.org/about.jsp

Cellular response to starvation ↗

Location: Cellular responses to stress

Stable identifier: R-MMU-9711097

Inferred from: Cellular response to starvation (Homo sapiens)

Amino acids regulate mTORC1



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

More details and caveats of the event inference in Reactome. For details on PANTHER see also: http://www.pantherdb.org/about.jsp

Unfolded Protein Response (UPR) 对


Location: Cellular responses to stress

Stable identifier: R-MMU-381119

Compartments: endoplasmic reticulum membrane, nucleoplasm, Golgi membrane, endoplasmic retic-

ulum lumen, cytosol

Inferred from: Unfolded Protein Response (UPR) (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.


More details and caveats of the event inference in Reactome. For details on PANTHER see also: http://www.pantherdb.org/about.jsp

Cellular response to mitochondrial stress 7

Location: Cellular responses to stress

Stable identifier: R-MMU-9840373

Inferred from: Cellular response to mitochondrial stress (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

 $\underline{More\ details\ and\ cave ats\ of\ the\ event\ inference\ in\ Reactome.}\ For\ details\ on\ PANTHER\ see\ also: \\ \underline{http://www.pantherdb.org/about.jsp}$

Table of Contents

Introduction	1
Cellular responses to stress	2
Cellular response to hypoxia	3
Cellular response to heat stress	4
Cellular Senescence	5
HSP90 chaperone cycle for steroid hormone receptors (SHR) in the presence of ligand	ϵ
Heme signaling	7
Cellular response to chemical stress	8
Cellular response to starvation	ç
Unfolded Protein Response (UPR)	10
Cellular response to mitochondrial stress	11
Table of Contents	12