

# Kcnq4 transports K+ from the cytosol to the extracellular region

Dallos, P., Furness, DN., May, B.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of <u>Creative Commons Attribution 4.0 International (CC BY 4.0)</u> <u>License</u>. For more information see our <u>license</u>.

26/09/2024

## Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

#### Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics, 18*, 142. 7
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. A
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res, 46*, D649-D655.
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, *14*, e1005968. *对*

This document contains 1 reaction (see Table of Contents)

### Kcnq4 transports K+ from the cytosol to the extracellular region 🛪

Stable identifier: R-MMU-9659557

Type: transition

#### Compartments: plasma membrane



Kcnq4 located on the basal membrane of outer hair cells (OHCs) (Kharkovets et al. 2000, Beisel et al. 2000, Kharkovets et al. 2006, Beisel et al. 2005, Winter et al. 2006) and inner hair cells (Kharkovets et al. 2000, Beisel et al. 2005) transports potassium ions along the concentration gradient from the cytosol to the extracellular region (Kharkovets et al. 2006). The resulting efflux of potassium is believed to be responsible for the I(K,n) current that plays a role in setting the resting potential of the cell and in repolarizing the cell (Oliver et al. 2003, Xu et al. 2007). Kcnq4 is located predominantly in the basal membrane of OHCs (Kharkovets et al. 2000, Winter et al. 2006) and absence of Kcnq4 causes increased depolarization and degeneration of OHCs (Kharkovets et al. 2006).

#### Literature references

- Fakler, B., Knipper, M., Oliver, D., Derst, C. (2003). Resting potential and submembrane calcium concentration of inner hair cells in the isolated mouse cochlea are set by KCNQ-type potassium channels. J. Neurosci., 23, 2141-9. 🛪
- El-Amraoui, A., Kharkovets, T., Safieddine, S., Jentsch, TJ., Hardelin, JP., Schweizer, M. et al. (2000). KCNQ4, a K+ channel mutated in a form of dominant deafness, is expressed in the inner ear and the central auditory pathway. *Proc. Natl. Acad. Sci. U.S.A.*, *97*, 4333-8.
- Christ, S., Walsh, EJ., Knipper, M., Bauer, K., Rohbock, K., McGee, J. et al. (2006). Thyroid hormone receptors TRalpha1 and TRbeta differentially regulate gene expression of Kcnq4 and prestin during final differentiation of outer hair cells. J. Cell. Sci., 119, 2975-84. *¬*
- Nelson, NC., Fritzsch, B., Delimont, DC., Beisel, KW. (2000). Longitudinal gradients of KCNQ4 expression in spiral ganglion and cochlear hair cells correlate with progressive hearing loss in DFNA2. *Brain Res. Mol. Brain Res.*, 82, 137-49. *¬*
- Kharkovets, T., Dedek, K., Maier, H., Nouvian, R., Moser, T., Leuwer, R. et al. (2006). Mice with altered KCNQ4 K+ channels implicate sensory outer hair cells in human progressive deafness. *EMBO J.*, *25*, 642-52.

#### **Editions**

2019-08-27	Authored, Edited	May, B.
2020-09-14	Reviewed	Furness, DN., Dallos, P.