

p-Y-Stat1,3,5 dimer enhances expression of Socs1

May, B., Touw, IP.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of <u>Creative Commons Attribution 4.0 International (CC BY 4.0)</u>
<u>License</u>. For more information see our <u>license</u>.

18/05/2024

https://reactome.org

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142.
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467.
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res, 46*, D649-D655.
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology, 14*, e1005968.

Reactome database release: 88

This document contains 1 reaction (see Table of Contents)


https://reactome.org Page 2

p-Y-Stat1,3,5 dimer enhances expression of Socs1 >

Stable identifier: R-MMU-9705455

Type: omitted

Compartments: nucleoplasm, cytosol



The Socs1 gene is transcribed to yield mRNA and the mRNA is translated to yield Socs1 protein. Stat1 dimers (Saito et al. 2000, Chan et al. 2014), Stat3 dimers (Naka et al. 1997), or Stat5 dimers (inferred from Stat1 and Stat3) presumed to be bound to the promoter of the Socs1 gene activate transcription of Socs1.

Literature references

Bhattacharya, D., Rickert, CG., Archambault, J., Vermi, W., White, JM., Lonardi, S. et al. (2014). Dysregulated STAT1-SOCS1 control of JAK2 promotes mammary luminal progenitor cell survival and drives ERα(+) tumorigenesis. *Cell Death Differ, 21*, 234-46. *¬*

Kajita, T., Kishimoto, T., Aono, A., Yoshizaki, K., Nishimoto, N., Matsumoto, T. et al. (1997). Structure and function of a new STAT-induced STAT inhibitor. *Nature*, 387, 924-9.

Editions

2020-10-18	Authored, Edited	May, B.
2020-12-12	Reviewed	Touw, IP.