

Tyrosine phosphorylation of Prlr

Goffin, V., Jupe, S.

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://reactome.org/licenses/).

06/05/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)

Reactome database release: 88


This document contains 1 reaction ([see Table of Contents](#))

Tyrosine phosphorylation of Prlr [↗](#)

Stable identifier: R-RNO-1369121

Type: transition

Compartments: plasma membrane, cytosol



The model for Prl-induced Prlr activation suggests that Jak2 phosphorylates Prlr on specific tyrosine residues. Consistent with this, Jak2 and Prlr are phosphorylated in response to activating ligand (Lebrun et al. 1994) and PRLR tyrosine phosphorylation is required for subsequent Stat signaling (Pezet et al. 1997). Though this evidence is consistent with a role for JAK2, it has not been formally demonstrated that JAK2 is the kinase responsible for PRLR tyrosine phosphorylation.

Literature references

Ferrag, F., Edery, M., Kelly, PA., Pezet, A. (1997). Tyrosine docking sites of the rat prolactin receptor required for association and activation of stat5. *J Biol Chem*, 272, 25043-50. [↗](#)

Goffin, V., Lebrun, JJ., Kelly, PA., Ullrich, A., Ali, S. (1995). A single phosphotyrosine residue of the prolactin receptor is responsible for activation of gene transcription. *Proc Natl Acad Sci U S A*, 92, 4031-5. [↗](#)

Editions

2011-06-13	Authored	Jupe, S.
2011-10-17	Edited	Jupe, S.
2011-11-08	Reviewed	Goffin, V.