

# **Regulation of TP53 Activity**

Regulation of TP53 Expression and Degradation	Regulation of TPS3 Activity through Phosphorylation	Regulation of TPS3 Activity through Activition	Regulation of TP53 Activity through Association with Co- factors	Regulation of TP53 Activity through Methylation	
				<b>r</b> eactome	

European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of <u>Creative Commons Attribution 4.0 International (CC BY 4.0)</u> <u>License</u>. For more information see our <u>license</u>.

This is just an excerpt of a full-length report for this pathway. To access the complete report, please download it at the <u>Reactome Textbook</u>.

16/05/2024

## Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

## Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics, 18,* 142. 7
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. A
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res, 46*, D649-D655. ↗
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, *14*, e1005968. *オ*

This document contains 6 pathways (see Table of Contents)

## **Regulation of TP53 Activity ↗**

#### Stable identifier: R-SSC-5633007

#### Inferred from: Regulation of TP53 Activity (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.


The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

## **Regulation of TP53 Expression and Degradation ↗**

Location: Regulation of TP53 Activity

Stable identifier: R-SSC-6806003

Inferred from: Regulation of TP53 Expression and Degradation (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.


The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

## Regulation of TP53 Activity through Phosphorylation 7

Location: Regulation of TP53 Activity

#### Stable identifier: R-SSC-6804756

Inferred from: Regulation of TP53 Activity through Phosphorylation (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.


The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

## Regulation of TP53 Activity through Association with Co-factors 7

Location: Regulation of TP53 Activity

#### Stable identifier: R-SSC-6804759

Inferred from: Regulation of TP53 Activity through Association with Co-factors (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.


The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

## Regulation of TP53 Activity through Methylation *オ*

Location: Regulation of TP53 Activity

#### Stable identifier: R-SSC-6804760

Inferred from: Regulation of TP53 Activity through Methylation (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.


The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

## Regulation of TP53 Activity through Acetylation 7

Location: Regulation of TP53 Activity

#### Stable identifier: R-SSC-6804758

Inferred from: Regulation of TP53 Activity through Acetylation (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

## **Table of Contents**

Introduction	1
暮 Regulation of TP53 Activity	2
暮 Regulation of TP53 Expression and Degradation	3
暮 Regulation of TP53 Activity through Phosphorylation	4
暮 Regulation of TP53 Activity through Association with Co-factors	5
暮 Regulation of TP53 Activity through Methylation	6
Transformation and TP53 Activity through Acetylation	7
Table of Contents	8