


Branched-chain amino acid catabolism



European Bioinformatics Institute, New York University Langone Medical Center, Ontario Institute for Cancer Research, Oregon Health and Science University.

The contents of this document may be freely copied and distributed in any media, provided the authors, plus the institutions, are credited, as stated under the terms of [Creative Commons Attribution 4.0 International \(CC BY 4.0\) License](https://creativecommons.org/licenses/by/4.0/). For more information see our [license](https://reactome.org/licenses/).

This is just an excerpt of a full-length report for this pathway. To access the complete report, please download it at the [Reactome Textbook](https://reactome.org/textbook/).

08/05/2024

Introduction

Reactome is open-source, open access, manually curated and peer-reviewed pathway database. Pathway annotations are authored by expert biologists, in collaboration with Reactome editorial staff and cross-referenced to many bioinformatics databases. A system of evidence tracking ensures that all assertions are backed up by the primary literature. Reactome is used by clinicians, geneticists, genomics researchers, and molecular biologists to interpret the results of high-throughput experimental studies, by bioinformaticians seeking to develop novel algorithms for mining knowledge from genomic studies, and by systems biologists building predictive models of normal and disease variant pathways.

The development of Reactome is supported by grants from the US National Institutes of Health (P41 HG003751), University of Toronto (CFREF Medicine by Design), European Union (EU STRP, EMI-CD), and the European Molecular Biology Laboratory (EBI Industry program).

Literature references

- Fabregat, A., Sidiropoulos, K., Viteri, G., Forner, O., Marin-Garcia, P., Arnau, V. et al. (2017). Reactome pathway analysis: a high-performance in-memory approach. *BMC bioinformatics*, 18, 142. [↗](#)
- Sidiropoulos, K., Viteri, G., Sevilla, C., Jupe, S., Webber, M., Orlic-Milacic, M. et al. (2017). Reactome enhanced pathway visualization. *Bioinformatics*, 33, 3461-3467. [↗](#)
- Fabregat, A., Jupe, S., Matthews, L., Sidiropoulos, K., Gillespie, M., Garapati, P. et al. (2018). The Reactome Pathway Knowledgebase. *Nucleic Acids Res*, 46, D649-D655. [↗](#)
- Fabregat, A., Korninger, F., Viteri, G., Sidiropoulos, K., Marin-Garcia, P., Ping, P. et al. (2018). Reactome graph database: Efficient access to complex pathway data. *PLoS computational biology*, 14, e1005968. [↗](#)


Reactome database release: 88

This document contains 1 pathway and 19 reactions ([see Table of Contents](#))

Branched-chain amino acid catabolism ↗

Stable identifier: R-XTR-70895

Inferred from: [Branched-chain amino acid catabolism \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

leu, ile, or val + alpha-ketoglutarate <=> a-ketoisocaproate, a-keto-b-methylvalerate, or a-ketoisovalerate + glutamate [BCAT1] ↗


Location: [Branched-chain amino acid catabolism](#)

Stable identifier: R-XTR-70723

Type: transition

Compartments: cytosol

Inferred from: [leu, ile, or val + alpha-ketoglutarate <=> a-ketoisocaproate, a-keto-b-methylvalerate, or a-ketoisovalerate + glutamate \[BCAT1\] \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

a-ketoisocaproate, a-keto-b-methylvalerate, or a-ketoisovalerate + glutamate <=> leu, ile, or val + alpha-ketoglutarate [BCAT1] ↗


Location: [Branched-chain amino acid catabolism](#)

Stable identifier: R-XTR-508189

Type: transition

Compartments: cytosol

Inferred from: [a-ketoisocaproate, a-keto-b-methylvalerate, or a-ketoisovalerate + glutamate <=> leu, ile, or val + alpha-ketoglutarate \[BCAT1\]](#) (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

leu, ile, or val + alpha-ketoglutarate <=> a-ketoisocaproate, a-keto-b-methylvalerate, or a-ketoisovalerate + glutamate [BCAT2] ↗


Location: [Branched-chain amino acid catabolism](#)

Stable identifier: R-XTR-70724

Type: transition

Compartments: mitochondrial matrix

Inferred from: [leu, ile, or val + alpha-ketoglutarate <=> a-ketoisocaproate, a-keto-b-methylvalerate, or a-ketoisovalerate + glutamate \[BCAT2\] \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Followed by: [BCKDH synthesizes BCAA-CoA from KIC, KMVA, KIV](#)

a-ketoisocaproate, a-keto-b-methylvalerate, or a-ketoisovalerate + glutamate <=> leu, ile, or val + alpha-ketoglutarate [BCAT2] ↗


Location: [Branched-chain amino acid catabolism](#)

Stable identifier: R-XTR-508179

Type: transition

Compartments: mitochondrial matrix

Inferred from: [a-ketoisocaproate, a-keto-b-methylvalerate, or a-ketoisovalerate + glutamate <=> leu, ile, or val + alpha-ketoglutarate \[BCAT2\]](#) (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

BCKDH synthesizes BCAA-CoA from KIC, KMVA, KIV ↗


Location: Branched-chain amino acid catabolism

Stable identifier: R-XTR-70713

Type: transition

Compartments: mitochondrial matrix

Inferred from: BCKDH synthesizes BCAA-CoA from KIC, KMVA, KIV (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: leu, ile, or val + alpha-ketoglutarate \rightleftharpoons a-ketoisocaproate, a-keto-b-methylvalerate, or a-ketoisovalerate + glutamate [BCAT2]

Followed by: isovaleryl-CoA + FAD \Rightarrow beta-methylcrotonyl-CoA + FADH₂, alpha-methylbutyryl-CoA + FAD \Rightarrow tiglyl-CoA + FADH₂, isobutyryl-CoA + FAD \Rightarrow methacrylyl-CoA + FADH₂

BCKDK phosphorylates BCKDH ↗


Location: Branched-chain amino acid catabolism

Stable identifier: R-XTR-5693148

Type: transition

Compartments: mitochondrial matrix

Inferred from: BCKDK phosphorylates BCKDH (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](http://www.pantherdb.org/about.jsp) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

isovaleryl-CoA + FAD => beta-methylcrotonyl-CoA + FADH2 ↗


Location: [Branched-chain amino acid catabolism](#)

Stable identifier: R-XTR-70745

Type: transition

Compartments: mitochondrial matrix

Inferred from: [isovaleryl-CoA + FAD => beta-methylcrotonyl-CoA + FADH2 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: [BCKDH synthesizes BCAA-CoA from KIC, KMVA, KIV](#)

Followed by: [beta-methylcrotonyl-CoA + ATP + CO2 <=> beta-methylglutaconyl-CoA + ADP + orthophosphate + H2O \[MCCA\]](#)

beta-methylcrotonyl-CoA + ATP + CO2 <=> beta-methylglutaconyl-CoA + ADP + orthophosphate + H2O [MCCA] ↗


Location: Branched-chain amino acid catabolism

Stable identifier: R-XTR-70773

Type: transition

Compartments: mitochondrial matrix

Inferred from: beta-methylcrotonyl-CoA + ATP + CO2 <=> beta-methylglutaconyl-CoA + ADP + orthophosphate + H2O [MCCA] (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](http://www.pantherdb.org/about.jsp) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: isovaleryl-CoA + FAD => beta-methylcrotonyl-CoA + FADH2

Followed by: beta-methylglutaconyl-CoA + H2O <=> beta-hydroxy-beta-methylglutaryl-CoA

beta-methylglutaconyl-CoA + ADP + orthophosphate + H2O <=> beta-methylcrotonyl-CoA + ATP + CO2 [MCCA] ↗


Location: Branched-chain amino acid catabolism

Stable identifier: R-XTR-508308

Type: transition

Compartments: mitochondrial matrix

Inferred from: beta-methylglutaconyl-CoA + ADP + orthophosphate + H2O <=> beta-methylcrotonyl-CoA + ATP + CO2 [MCCA] (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](http://www.pantherdb.org/about.jsp) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

beta-methylglutaconyl-CoA + H2O <=> beta-hydroxy-beta-methylglutaryl-CoA ↗


Location: [Branched-chain amino acid catabolism](#)

Stable identifier: R-XTR-70785

Type: transition

Compartments: mitochondrial matrix

Inferred from: [beta-methylglutaconyl-CoA + H2O <=> beta-hydroxy-beta-methylglutaryl-CoA \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: [beta-methylcrotonyl-CoA + ATP + CO2 <=> beta-methylglutaconyl-CoA + ADP + orthophosphate + H2O \[MCCA\]](#)

alpha-methylbutyryl-CoA + FAD => tiglyl-CoA + FADH2 ↗


Location: Branched-chain amino acid catabolism

Stable identifier: R-XTR-70800

Type: transition

Compartments: mitochondrial matrix

Inferred from: alpha-methylbutyryl-CoA + FAD => tiglyl-CoA + FADH2 (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](http://www.pantherdb.org/about.jsp) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: BCKDH synthesizes BCAA-CoA from KIC, KMVA, KIV

alpha-methyl-beta-hydroxybutyryl-CoA + NAD+ <=> alpha-methylacetoacetyl-CoA + NADH + H+ ↗


Location: Branched-chain amino acid catabolism

Stable identifier: R-XTR-70837

Type: transition

Compartments: mitochondrial matrix

Inferred from: alpha-methyl-beta-hydroxybutyryl-CoA + NAD+ <=> alpha-methylacetoacetyl-CoA + NADH + H+ (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](https://www.reactome.org/docs/inferring_events.html) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Followed by: alpha-methyl-acetoacetyl-CoA + CoA => propionyl-CoA + acetyl-CoA

alpha-methylacetoacetyl-CoA + NADH + H+ <=> alpha-methyl-beta-hydroxybutyryl-CoA + NAD+ ↗


Location: Branched-chain amino acid catabolism

Stable identifier: R-XTR-508369

Type: transition

Compartments: mitochondrial matrix

Inferred from: alpha-methylacetoacetyl-CoA + NADH + H+ <=> alpha-methyl-beta-hydroxybutyryl-CoA + NAD+ (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](https://www.reactome.org) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

alpha-methyl-acetoacetyl-CoA + CoA => propionyl-CoA + acetyl-CoA ↗


Location: Branched-chain amino acid catabolism

Stable identifier: R-XTR-70844

Type: transition

Compartments: mitochondrial matrix

Inferred from: alpha-methyl-acetoacetyl-CoA + CoA => propionyl-CoA + acetyl-CoA (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](http://www.pantherdb.org/about.jsp) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: alpha-methyl-beta-hydroxybutyryl-CoA + NAD+ <=> alpha-methylacetoacetyl-CoA + NADH + H+

isobutyryl-CoA + FAD => methacrylyl-CoA + FADH2 ↗


Location: [Branched-chain amino acid catabolism](#)

Stable identifier: R-XTR-70859

Type: transition

Compartments: mitochondrial matrix

Inferred from: [isobutyryl-CoA + FAD => methacrylyl-CoA + FADH2 \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](#) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: [BCKDH synthesizes BCAA-CoA from KIC, KMVA, KIV](#)

Followed by: [ECHS1 hydrates methacrylyl-CoA](#)

ECCHS1 hydrates methacrylyl-CoA ↗


Location: Branched-chain amino acid catabolism

Stable identifier: R-XTR-70870

Type: transition

Compartments: mitochondrial matrix

Inferred from: ECCHS1 hydrates methacrylyl-CoA (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](http://www.pantherdb.org/about.jsp) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: isobutyryl-CoA + FAD => methacrylyl-CoA + FADH₂

beta-hydroxyisobutyrate + NAD+ <=> methylmalonyl semialdehyde + NADH + H+ ↗


Location: Branched-chain amino acid catabolism

Stable identifier: R-XTR-70885

Type: transition

Compartments: mitochondrial matrix

Inferred from: beta-hydroxyisobutyrate + NAD+ <=> methylmalonyl semialdehyde + NADH + H+ (Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](http://www.pantherdb.org/about.jsp) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Followed by: methylmalonate semialdehyde + NAD+ + CoA => propionyl-CoA + CO₂ + NADH + H+

methylmalonyl semialdehyde + NADH + H+ <=> beta-hydroxyisobutyrate + NAD+ ↗


Location: [Branched-chain amino acid catabolism](#)

Stable identifier: R-XTR-508473

Type: transition

Compartments: mitochondrial matrix

Inferred from: [methylmalonyl semialdehyde + NADH + H+ <=> beta-hydroxyisobutyrate + NAD+ \(Homo sapiens\)](#)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](http://www.pantherdb.org/about.jsp) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

methyImalonate semialdehyde + NAD⁺ + CoA => propionyl-CoA + CO₂ + NADH + H⁺ ↗


Location: Branched-chain amino acid catabolism

Stable identifier: R-XTR-70893

Type: transition

Compartments: mitochondrial matrix

Inferred from: methyImalonate semialdehyde + NAD⁺ + CoA => propionyl-CoA + CO₂ + NADH + H⁺
(Homo sapiens)



This event has been computationally inferred from an event that has been demonstrated in another species.

The inference is based on the homology mapping from PANTHER. Briefly, reactions for which all involved PhysicalEntities (in input, output and catalyst) have a mapped orthologue/paralogue (for complexes at least 75% of components must have a mapping) are inferred to the other species. High level events are also inferred for these events to allow for easier navigation.

[More details and caveats of the event inference in Reactome.](http://www.pantherdb.org/about.jsp) For details on PANTHER see also: <http://www.pantherdb.org/about.jsp>

Preceded by: beta-hydroxyisobutyrate + NAD⁺ <=> methylmalonyl semialdehyde + NADH + H⁺

Table of Contents

Introduction	1
☞ Branched-chain amino acid catabolism	2
☞ leu, ile, or val + alpha-ketoglutarate \rightleftharpoons a-ketoisocaproate, a-keto-b-methylvalerate, or a-ketoisovalerate + glutamate [BCAT1]	3
☞ a-ketoisocaproate, a-keto-b-methylvalerate, or a-ketoisovalerate + glutamate \rightleftharpoons leu, ile, or val + alpha-ketoglutarate [BCAT1]	4
☞ leu, ile, or val + alpha-ketoglutarate \rightleftharpoons a-ketoisocaproate, a-keto-b-methylvalerate, or a-ketoisovalerate + glutamate [BCAT2]	5
☞ a-ketoisocaproate, a-keto-b-methylvalerate, or a-ketoisovalerate + glutamate \rightleftharpoons leu, ile, or val + alpha-ketoglutarate [BCAT2]	6
☞ BCKDH synthesizes BCAA-CoA from KIC, KMVA, KIV	7
☞ BCKDK phosphorylates BCKDH	8
☞ isovaleryl-CoA + FAD \Rightarrow beta-methylcrotonyl-CoA + FADH ₂	9
☞ beta-methylcrotonyl-CoA + ATP + CO ₂ \rightleftharpoons beta-methylglutaconyl-CoA + ADP + orthophosphate + H ₂ O [MCCA]	10
☞ beta-methylglutaconyl-CoA + ADP + orthophosphate + H ₂ O \rightleftharpoons beta-methylcrotonyl-CoA + ATP + CO ₂ [MCCA]	11
☞ beta-methylglutaconyl-CoA + H ₂ O \rightleftharpoons beta-hydroxy-beta-methylglutaryl-CoA	12
☞ alpha-methylbutyryl-CoA + FAD \Rightarrow tiglyl-CoA + FADH ₂	13
☞ alpha-methyl-beta-hydroxybutyryl-CoA + NAD ⁺ \rightleftharpoons alpha-methylacetoacetyl-CoA + NADH + H ⁺	14
☞ alpha-methylacetoacetyl-CoA + NADH + H ⁺ \rightleftharpoons alpha-methyl-beta-hydroxybutyryl-CoA + NAD ⁺	15
☞ alpha-methyl-acetoacetyl-CoA + CoA \Rightarrow propionyl-CoA + acetyl-CoA	16
☞ isobutyryl-CoA + FAD \Rightarrow methacrylyl-CoA + FADH ₂	17
☞ ECHS1 hydrates methacrylyl-CoA	18
☞ beta-hydroxyisobutyrate + NAD ⁺ \rightleftharpoons methylmalonyl semialdehyde + NADH + H ⁺	19
☞ methylmalonyl semialdehyde + NADH + H ⁺ \rightleftharpoons beta-hydroxyisobutyrate + NAD ⁺	20
☞ methylmalonate semialdehyde + NAD ⁺ + CoA \Rightarrow propionyl-CoA + CO ₂ + NADH + H ⁺	21
Table of Contents	22